

Class 9: 2/17/14 Section 3.3. I

We start with the continuity of compositions of functions.

Thm If  $g(x)$  is continuous at  $x=c$ , with  $g(c)=L$ , and  $f(x)$  is continuous at  $x=L$ , then  $(f \circ g)(x)$  is continuous at  $x=c$ , and

$$\begin{aligned}\lim_{x \rightarrow c} (f \circ g)(x) &= \lim_{x \rightarrow c} f(g(x)) = f\left(\lim_{x \rightarrow c} g(x)\right) \\ &= f(g(c)) = f(L).\end{aligned}$$

Notes ① Typically, one must work from the inside out when establishing continuity of a composition at a pt  $x=c$ .

ex. Where is  $f(x) = \sin(\ln x)$  continuous?

Here the inside function  $\ln x$  is cont on  $(0, \infty)$ , with range all  $\mathbb{R}$ . And since  $\sin x$  is cont. on  $\mathbb{R}$ ,  $f(x)$  is cont on  $(0, \infty)$ .

ex.  $g(x) = \ln(\sin x)$ . Where is  $g(x)$  cont?

Here, the inside function  $\sin x$  is cont on all of  $\mathbb{R}$ , but its range is  $[-1, 1]$ .

The outside function is cont. only on  $(0, \infty)$

Hence only those input values whose range is in  $(0, 1] = [-1, 1] \cap (0, \infty)$  are where  $g(x)$  is continuous. Where is  $\sin x \in (0, 1]$ ?

$g(x)$  is continuous on the domain

$$\begin{aligned} & \{x \in \mathbb{R} \mid \sin x \in (0, 1]\} \\ &= \{x \in \mathbb{R} \mid 0 < x \bmod 2\pi < \pi\} \end{aligned}$$

Think about this one.

---

ex. (7ed pg 107)

Determine where  $h(x) = \frac{1}{1+2x^{1/3}}$  is continuous.

Note: The book uses  $g(x) = x^{1/3}$  and  $f(x) = \frac{1}{1+2x}$  to write  $h(x) = f(g(x)) = (f \circ g)(x)$ . The result is  $h(x)$  is continuous on  $\{x \in \mathbb{R} \mid x \neq -\frac{1}{2}\}$ .

Here, we split  $h(x)$  using  $f(x) = \frac{1}{x}$  and  $g(x) = 1+2x^{1/3}$  so that  $h(x) = f(g(x)) = f(1+2x^{1/3}) = \frac{1}{1+2x^{1/3}}$ .

Here,  $g(x) = 1+2x^{1/3}$  is continuous everywhere, and its range is all of  $\mathbb{R}$ . But  $f(x)$  is not defined when  $x=0$ , so we need to rule out all input values for  $g(x)$  where the output value is 0: rule out all  $x \in \mathbb{R}$  where  $g(x) = 0$ .

$g(x) = 0$  when  $1+2x^{1/3} = 0$ , or  $x^{1/3} = -\frac{1}{2}$ , or  $x = -\frac{1}{8}$ .

Hence  $h(x) = f(g(x)) = \frac{1}{1+2x^{1/3}}$  is continuous on the domain  $\{x \in \mathbb{R} \mid x \neq -\frac{1}{8}\} = \mathbb{R} \setminus \{-\frac{1}{8}\} = (-\infty, -\frac{1}{8}) \cup (-\frac{1}{8}, \infty)$ .

Recall a sequence  $\{a_n\}_{n=1}^{\infty}$  has a limit

$L$ , and we say  $\lim_{n \rightarrow \infty} a_n = L$ , if

for every  $\varepsilon > 0$ , there is an  $N \in \mathbb{N}$  so that

$$|a_n - L| < \varepsilon \text{ whenever } n > N.$$

We used this to show  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ . Given

any  $\varepsilon > 0$ , choose  $N = \frac{1}{\varepsilon}$ . (can you see this?)

---

This also works for functions of a continuous variable  $x \in \mathbb{R}$ :

A function  $f(x)$  has a limit  $L$  at infinity,


denoted  $\lim_{x \rightarrow \infty} f(x) = L$ , if for every  $\varepsilon > 0$

there is an  $x_0 \in \mathbb{R}$ , so that

$$|f(x) - L| < \varepsilon \text{ whenever } x > x_0$$

Recognize that this is a formal, precise definition of a limit at infinity.

Visually,




Can be used to show

$$\lim_{x \rightarrow \infty} \frac{1}{x} = 0. \text{ Given } \epsilon > 0, \text{ we seek}$$

a pt  $x_0 \in \mathbb{R}$  when function  $f(x)$  will always lie between 0 and  $\epsilon$  ever afterward.

Given an  $\epsilon > 0$ , we can calculate  $x_0$ :



Want

$$|f(x) - L| < \epsilon \text{ whenever } x > x_0.$$

$$\left| \frac{1}{x} - 0 \right| < \epsilon \Rightarrow \frac{1}{x} < \epsilon. \text{ This will be true when } x > \frac{1}{\epsilon}.$$

So choose  $x_0 = \frac{1}{\epsilon}$ . Then def of limit holds.

ex. Show  $\lim_{x \rightarrow \infty} \frac{1}{x^n} = 0$ , for  $n \in \mathbb{N}$ .


Method 1: By definition

want  $|f(x) - L| < \epsilon$

when  $x > x_0$ .

Choose  $L = 0$ .

$$|f(x) - L| = \left| \frac{1}{x^n} - 0 \right| = \frac{1}{x^n}$$



and  $\frac{1}{x^n} < \epsilon$ , precisely when  $\sqrt[n]{\frac{1}{\epsilon}} < x$ .

choose  $x_0 = \sqrt[n]{\frac{1}{\epsilon}}$ . Then def holds, and

$$\lim_{x \rightarrow \infty} \frac{1}{x^n} = 0.$$

Method 2: Use limit laws: (and previous result)

$$\lim_{x \rightarrow \infty} \frac{1}{x^n} = \lim_{x \rightarrow \infty} \left( \frac{1}{x} \right)^n = \lim_{x \rightarrow \infty} \overbrace{\left( \frac{1}{x} \right) \left( \frac{1}{x} \right) \dots \left( \frac{1}{x} \right)}^{n \text{ times}}$$

Product Rule  
for limits  $\left( \lim_{x \rightarrow \infty} \frac{1}{x} \right) \left( \lim_{x \rightarrow \infty} \frac{1}{x} \right) \dots \left( \lim_{x \rightarrow \infty} \frac{1}{x} \right)$


as long as each of these exists.

Then do, and

$$= 0 \cdot 0 \cdot \dots \cdot 0 = 0.$$

ex. Calculate  $\lim_{x \rightarrow \infty} x$ , if it exists.

Suppose you choose any  
real number  $L \geq 0$ .  
as a possible limit



Then for any choice of  $\epsilon > 0$ , once the function  
~~the~~ input values get larger than  $L + \epsilon$ ,  
 $f(x) = x$  will leave the band and never  
return. Hence there is no real number  $L$   
that can serve as the limit.  
Hence limit does not exist.

Note: This kind of nonexisting limit is  
particular. We say  $\lim_{x \rightarrow \infty} x = \infty$ .


This is different from  $\lim_{x \rightarrow \infty} \sin x$ , which  
also does not exist.

See box pg 95 and next example.

ex. Calculate  $\lim_{x \rightarrow \infty} \sin x$ , if it exists.

It would make no sense, if  $L$  exists,

that  $L > 1$  or  $L < -1$  (why?).




So choose some  $L \in [-1, 1]$  to possibly serve as the limit. Choose an  $\epsilon > 0$  small enough, like in the picture. Can you see that there will be no  $x_0 \in \mathbb{R}$  where for all  $x > x_0$ ,  $|\sin x - L| < \epsilon$  ??

$\lim_{x \rightarrow \infty} \sin x$  does not exist.

ex. Show  $\lim_{x \rightarrow \infty} e^{-x} = 0$ .

Like  $\frac{1}{x}$ , we seek  $\epsilon$  pt  $x_0 \in \mathbb{R}$  where when  $x > x_0$ ,

$$|f(x) - L| = |e^{-x} - 0| = e^{-x} < \epsilon.$$





Note: ① Polynomials (of positive degree) never have limits at infinity or  $-\infty$  (why not?)

② Rational functions sometimes do end sometimes do not. It turns out there is an easy way to tell!!

Thm For  $f(x) = \frac{p(x)}{q(x)}$  rational, we have

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{p(x)}{q(x)} = \begin{cases} 0 & \text{if } \deg(p) < \deg(q) \\ L & \text{if } \deg(p) = \deg(q) \\ \infty & \text{if } \deg(p) > \deg(q) \end{cases}$$

where  $L$  is a real number. (what is  $L$ ?)

Note: The proof is not difficult and kinda interesting. The next example should give you an idea of how it would work.

ex. Show  $\lim_{x \rightarrow \infty} \frac{3x^2 - 4x + 5}{7x^3 + 3x^2 + 6} = 0.$

Strategy: Use the limit laws and a little manipulation of the rational function to calculate the limit directly.

Solution: The previous theorem establishes this

$$\text{since } \deg(p) = \deg(3x^2 - 4x + 5) = 2.$$

$$\text{and } \deg(q) = \deg(7x^3 + 3x^2 + 6) = 3.$$

Since  $f(x) = \frac{3x^2 - 4x + 5}{7x^3 + 3x^2 + 6}$  is unknown on

$(0, \infty)$  (why?), we know it is possible

for the limit to exist. We cannot apply the Quotient Law for limits

directly since  $\lim_{x \rightarrow \infty} p(x) = \lim_{x \rightarrow \infty} 3x^2 - 4x + 5$

does not exist.

ex (cont'd)

But, using a common trick in math; multiplied by a "clever form of one", we can manipulate  $f(x)$  so that we can use the Quotient Rule for limits:

$$1 = \left( \frac{1}{\frac{1}{x^3}} \right) \text{ is continuous on } (0, \infty), \text{ and}$$

thus so is

$$\frac{(3x^2 - 4x + 5) \left( \frac{1}{x^3} \right)}{(7x^3 + 3x^2 + 6) \left( \frac{1}{x^3} \right)} = \frac{\frac{3}{x} - \frac{4}{x^2} + \frac{5}{x^3}}{7 + \frac{3}{x} + \frac{6}{x^3}}$$

since it is the product of 2 continuous functions on  $(0, \infty)$ . Hence

$$\begin{aligned} \lim_{x \rightarrow \infty} f(x) &= \lim_{x \rightarrow \infty} \frac{3x^2 - 4x + 5}{7x^3 + 3x^2 + 6} = \lim_{x \rightarrow \infty} \frac{(3x^2 - 4x + 5) \left( \frac{1}{x^3} \right)}{(7x^3 + 3x^2 + 6) \left( \frac{1}{x^3} \right)} \\ &= \lim_{x \rightarrow \infty} \frac{\frac{3}{x} - \frac{4}{x^2} + \frac{5}{x^3}}{7 + \frac{3}{x} + \frac{6}{x^3}} \end{aligned}$$

ex. (cont'd) Now for  $A(x) = \frac{p(x)}{q(x)} = \frac{\frac{3}{x} - \frac{4}{x^2} + \frac{5}{x^3}}{7 + \frac{2}{x} + \frac{6}{x^3}}$

Now, since  $\lim_{x \rightarrow \infty} p(x) = \lim_{x \rightarrow \infty} \left( \frac{3}{x} - \frac{4}{x^2} + \frac{5}{x^3} \right)$

~~for~~  
~~for~~  
~~for~~

$$\begin{aligned} & \stackrel{\text{sum}}{\text{Rule}} \lim_{x \rightarrow \infty} \frac{3}{x} + \lim_{x \rightarrow \infty} \frac{-4}{x^2} + \lim_{x \rightarrow \infty} \frac{5}{x^3} \\ & \stackrel{\text{const}}{\text{mult}} \text{Rule} 3 \left( \lim_{x \rightarrow \infty} \frac{1}{x} \right) - 4 \left( \lim_{x \rightarrow \infty} \frac{1}{x^2} \right) + 5 \left( \lim_{x \rightarrow \infty} \frac{1}{x^3} \right) \\ & = 3(0) - 4(0) + 5(0) = 0 \end{aligned}$$

and  $\lim_{x \rightarrow \infty} q(x) = \lim_{x \rightarrow \infty} \left( 7 - \frac{2}{x} + \frac{6}{x^3} \right) = 7 - 2(0) + 6(0)$

by the same reasoning, we get

$$\lim_{x \rightarrow \infty} A(x) = \lim_{x \rightarrow \infty} \frac{p(x)}{q(x)} = \frac{\lim_{x \rightarrow \infty} p(x)}{\lim_{x \rightarrow \infty} q(x)} = \frac{0}{7} = 0$$

by the Quotient Rule for limits.

Can you see that this "trick" (multiplication) by a "clever form of 1", can be used in the proof of the theorem?