

# JOHNS HOPKINS MATH TOURNAMENT 2019

## Individual Round: Algebra

*February 9, 2019*

### Instructions

- **DO NOT TURN OVER THIS PAPER UNTIL TOLD TO DO SO.**
- This test contains 10 questions to be solved individually in 60 minutes.
- All answers will be integers.
- Only answers written on the appropriate area on the answer sheet will be considered for grading.
- Problems are weighted relative to their difficulty, determined by the number of students who solve each problem.
- No translators, books, notes, slide rules, calculators, abaci, or other computational aids are permitted. Similarly, graph paper, rulers, protractors, compasses, and other drawing aids are not permitted.
- If you believe the test contains an error, immediately tell your proctor and if necessary, report the error to the front desk after the end of your exam.
- Good luck!

- The sum of the squares of two numbers is 2019 and the product of the two numbers is 95. What is the sum of the two numbers?
- How many ordered pairs of integers  $(a, b)$  exist such that  $1/a + 1/b = 1/4$ ?
- How many natural numbers less than 2019 are there such that its remainder when divided by 2 is 1, when divided by 3 is 2, when divided by 4 is 3, and when divided by 5 is 4?

- Let  $P(x)$  be a polynomial with real coefficients where the sum of the coefficients is equal to 2019. Also,  $P(x)$  satisfies

$$P(-x) = -P(x)$$

The remainder,  $Q(x)$ , obtained by dividing  $P(x)$  by  $x^3 - x$  has the form  $px^2 + qx + r$ , where  $p, q$ , and  $r$  are constants. Find  $p + q + r$ .

- The given polynomial  $P(X)$  has leading coefficient 1 and satisfies the functional equation below:

$$(X + 1)P(X) = (X - 10)P(X + 1)$$

Compute  $P(5)$ .

- $\binom{1000}{0} - \binom{1000}{2} + \binom{1000}{4} - \dots + \binom{1000}{1000} = 2^A$ . Find  $A$ .
- Given the quadratic equation  $ax^2 - bx + c = 0$ , where  $a, b, c \in \mathbb{R}$ , find the coefficients  $a, b, c$  such that the equation has the roots  $a, b$  and discriminant  $c$ . Compute  $\frac{4c}{ab}$ .
- The equation below has only one real solution of the form  $a/b$  where  $a$  and  $b$  are coprime. Find  $a + b$ .

$$x^3 + (x + 1)^3 + (x + 2)^3 + (x + 3)^3 = 0$$

- Given that the equation  $(1 + x + x^2 + x^3 + \dots + x^{17})^2 - x^{17} = 0$  has 34 complex roots of the form  $z_k = r_k[\cos(2\pi a_k) + i \sin(2\pi a_k)]$ ,  $k = 1, 2, 3, \dots, 34$ , with  $0 < a_1 \leq a_2 \leq a_3 \leq \dots \leq a_{34} < 1$  and  $r_k > 0$ . Find  $a_1 + a_2 + a_3 + a_4 + a_5$ . Given the answer is in the form  $\frac{\alpha}{\beta}$ , compute  $\alpha + \beta$ .
- The roots,  $a, b, c$ , of the equation  $x^3 - 4x^2 + 5x - 19/10 = 0$  are real and can form the sides of a triangle. Given the area of the triangle has form  $\sqrt{q}/p$  where  $p$  is an integer and  $\sqrt{q}$  is in simplest radical form, find  $p + q$ .