EXAMPLE: THE WRONSKIAN DETERMINANT OF A SECOND-ORDER, LINEAR HOMOGENEOUS DIFFERENTIAL EQUATION

110.302 DIFFERENTIAL EQUATIONS PROFESSOR RICHARD BROWN

Given a second order, linear, homogeneous differential equation

$$y'' + p(t)y' + q(t)y = 0,$$

where both p(t) and q(t) are continuous on some open t-interval I, and two solutions $y_1(t)$ and $y_2(t)$, one can form a fundamental set of solutions as the linear combination of these two

$$y(t) = c_1 y_1(t) + c_2 y_2(t)$$

ONLY under the condition that the Wronskian determinant

$$W(y_1, y_2)(t) = \begin{vmatrix} y_1(t) & y_2(t) \\ y'_1(t) & y'_2(t) \end{vmatrix} \neq 0$$

for all $t \in I$. This condition implies that the two differentiable functions $y_1(t)$ and $y_2(t)$ are independent; that there does not exist two constants $k_1, k_2 \in \mathbb{R}$, not both equal to 0, where $k_1y_1(t) + k_2y_2(t) = 0$ on I. Then we know that ALL solutions to the ODE will be a linear combination of $y_1(t)$ and $y_2(t)$.

We can actually take this further. It turns out that in this situation, and on the interval I, either of two things can happen:

- The Wronskian is always 0 on I (we say $W(y_1, y_1)$ is identically 0 on I, or $W(y_1, y_2) \equiv 0$ on I), or
- the Wronskian is NEVER 0 on I.

Why? Because of the following:

Theorem. If $y_1(t)$ and $y_2(t)$ are two solutions to the ODE y'' + p(t)y' + q(t)y = 0, where p(t) and q(t) are continuous on some open t-interval I, then

$$W(y_1, y_2)(t) = Ce^{-\int p(t) dt}$$

where C depends on the choice of y_1 and y_2 , but not on t.

First, some notes:

- If $y_1(t)$ and $y_2(t)$ are linearly dependent, then C = 0.
- If $y_1(t)$ and $y_2(t)$ are linearly independent, then $W(y_1, y_2) \neq 0$ on ALL of I.
- Linear independence and non-zero Wronskian are the same thing for solutions to these ODEs.

Proof. Since y_1 and y_1 both solve the ODE, we have

$$y_1'' + p(t)y_1' + q(t)y_1 = 0$$

$$y_2'' + p(t)y_2' + q(t)y_2 = 0.$$

Here, multiply the first equation by $-y_2$ and the second by y_1 and then add them together (this will eliminate the coefficient q(t))

$$\underbrace{(y_1y_2''-y_2y_1'')}_{W'(y_1,y_2)} + p(t)\underbrace{(y_1y_2'-y_2y_1')}_{W(y_1,y_2)} = 0,$$

Which leads to a first order differential equation whose variable IS the Wronskian determinant itself (really, the Wronskian is a function of t)

$$W' + p(t)W = 0.$$

This first order ODE is both linear and separable, and by separation of variables, we get

$$\frac{W'}{W} = -p(t) \implies \ln|W| = -\int p(t) dt + K \implies Ce^{-\int p(t) dt}.$$

Hence by the notes above just before the proof, either C = 0, and the Wronskian is always 0, and the two solutions are linearly dependent, or $C \neq 0$, and the Wronskian is NEVER 0, and the two solutions are linearly independent.