

110.302 Lecture 25: 10/30/15 I

New question: What if the eigenvalues (solutions to the characteristic eqn of $A_{2 \times 2}$ in $\vec{x}' = A\vec{x}$ are not real?

Then they are complex (the discriminant $b^2 - 4ac$ of the quadratic formula used to solve the char. eqn is < 0).

Reqs must be complex conjugates (why?)

How to use them?

Let's play the same game for constructing solutions to $\vec{x}' = A\vec{x}$ using eigenvalues/eigenvector pairs:

For $\vec{X}' = A_{2 \times 2} \vec{X}$, suppose $r_1 \neq r_2$ are two distinct solns to char. eqn of A and we calculate eigenvectors $\vec{v}_1, \vec{v}_2$ respectively to r_1, r_2 . Then general soln is

$$\vec{X}(t) = c_1 \vec{v}_1 e^{r_1 t} + c_2 \vec{v}_2 e^{r_2 t}$$

We try this with complex r :

ex. $\vec{X}' = \begin{bmatrix} 0 & 1 \\ -2 & -2 \end{bmatrix} \vec{X}$. char. eqn is $r^2 + 2r + 2 = 0$,
solved by $r = -1 \pm i$

Leave them as $r_1 = -1 + i$, $r_2 = -1 - i$ and solve

for $\vec{v}_1$: $A\vec{v} = r\vec{v}$

$$\begin{bmatrix} 0 & 1 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = (-1 + i) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, \text{ or}$$

$$v_2 = -v_1 + i v_1$$

$$-2v_1 - 2v_2 = -v_2 + i v_2$$

We can substitute (1) into (2) and

simplify to get $-2i v_1 = -2i v_1$, solved by any choice of v . For example, choose $v_1 = 1$. Then $v_2 = (-1+i)$ and

$$v_1 = -1+i, \vec{v}_1 = \begin{bmatrix} 1 \\ -1+i \end{bmatrix}$$

forms an "eigenvalue/eigenvector" pair

Notes ① This is not quite accurate since the definition of eigenvector is that of a vector whose direction does not change upon mult by a matrix. Without real eigenvalues, there are no real eigenvectors! But the term "complex eigenvector" is a commonly used one.

Notes cont'd.

② No other eigenvalue/eigenvector pair is
 $r_2 = -1 - i$, $\vec{v}_2 = \begin{bmatrix} 1 \\ -1-i \end{bmatrix}$. (check this!)

③ Rewrite $\vec{v}_1 = \begin{bmatrix} 1 \\ -1+i \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} i = \vec{a} + i\vec{b}$

Then along with $r_1 = -1 + i = \lambda + i\mu$
 we can attempt to form solutions.

General idea for a Method for constructing solutions?

- Given $r_1 = \lambda + i\mu$, $\vec{v}_1 = \vec{a} + i\vec{b}$
 $r_2 = \lambda - i\mu$, $\vec{v}_2 = \vec{a} - i\vec{b}$,

Create a "complex" solution in the normal way:

$$\begin{aligned} \vec{x}(t) &= \vec{v}_1 e^{r_1 t} = (\vec{a} + i\vec{b}) \underbrace{e^{\lambda t} (\cos \mu t + i \sin \mu t)}_{\text{Euler formula for } e^{i\mu t}} \\ &= e^{\lambda t} (\vec{a} \cos \mu t - \vec{b} \sin \mu t) \\ &\quad + i e^{\lambda t} (\vec{a} \sin \mu t + \vec{b} \cos \mu t) \end{aligned}$$

V

Hence we can write $\vec{x}(t) = \vec{u}(t) + i\vec{w}(t)$, where

$$\vec{u}(t) = e^{\lambda t} (\vec{a} \cos \mu t - \vec{b} \sin \mu t)$$

$$\vec{w}(t) = e^{\lambda t} (\vec{a} \sin \mu t + \vec{b} \cos \mu t)$$

Notes ① These are 2 real-valued functions which each solve the ODE $\vec{x}' = A\vec{x}$ (check this!)

② They are independent (check the Wronskian)

③ The general solution to $\vec{x}' = A\vec{x}$ when eigenvalues $\gamma = \lambda \pm i\mu$ are complex and with eigenvectors $\vec{v} = \vec{a} \pm i\vec{b}$ is

$$\vec{x}(t) = c_1 \vec{u}(t) + c_2 \vec{w}(t)$$

Back to example: $\vec{x}' = \begin{bmatrix} 0 & 1 \\ -2 & -2 \end{bmatrix} \vec{x}$.

Here $\gamma_1 = \lambda + i\mu = -1 + i$, $\vec{v}_1 = \vec{a} + i\vec{b} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} + i \begin{bmatrix} 0 \\ 1 \end{bmatrix}$,

so

$$\vec{x}(t) = c_1 e^{-t} \left(\begin{bmatrix} 1 \\ -1 \end{bmatrix} \cos t - \begin{bmatrix} 0 \\ 1 \end{bmatrix} \sin t \right) + c_2 e^{-t} \left(\begin{bmatrix} 1 \\ -1 \end{bmatrix} \sin t + \begin{bmatrix} 0 \\ 1 \end{bmatrix} \cos t \right)$$