

110.302 Lecture 28: 11/06/15

I

Now consider the system

$$\begin{aligned} \dot{x}_1 &= -2x_1 + x_2 \\ \dot{x}_2 &= -2x_2 \end{aligned} \quad \text{or} \quad \vec{x}' = \begin{bmatrix} -2 & 1 \\ 0 & -2 \end{bmatrix} \vec{x}$$

Here eigenvalues of $A = \begin{bmatrix} -2 & 1 \\ 0 & -2 \end{bmatrix}$ satisfy $r^2 + 4r + 4 = 0$
or $r_1 = -2 = r_2$.

Eigenvectors satisfy $A\vec{v} = -2\vec{v}$:

$$\begin{aligned} -2v_1 + v_2 &= -2v_1 \\ -2v_2 &= -2v_2 \end{aligned} \quad \left. \vphantom{\begin{aligned} -2v_1 + v_2 &= -2v_1 \\ -2v_2 &= -2v_2 \end{aligned}} \right\} \begin{aligned} v_2 &= 0 \\ v_1 &= \text{anything} \end{aligned}$$

All eigenvectors look like $\begin{bmatrix} * \\ 0 \end{bmatrix}$. Hence there is only 1 lin. indep. eigenvector. Choose $*=1$.

For $r = -2$, $\vec{v} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$. and 1 soln is $\vec{x}_1(t) = \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^{-2t}$.

How to find another one?

Q: What did we do for the case of a 2^{nd} -order lin. homogeneous ODE with constant coefficients when there was only one root to the char. eqn?

A: Two indep solutions were e^{-2t} and te^{-2t} .

Lets try this here:

Create a guess for a solution:

$$\vec{x}^{(2)}(t) = \vec{w} t e^{-2t} \quad \text{for } \vec{w} \text{ a 2-vector}$$

For this to be a solution, it must "fit" the

$$\text{ODE: } \vec{x}' = A\vec{x}$$

For this example, we set

$$\frac{d}{dt} [\vec{w} t e^{-2t}] = \begin{bmatrix} -2 & 1 \\ 0 & -2 \end{bmatrix} \cancel{e^{-2t}} (\vec{w} t e^{-2t})$$

$$\vec{w} e^{-2t} - 2\vec{w} t e^{-2t} = A\vec{w} t e^{-2t}$$

$$\vec{w} - 2\vec{w} t = A\vec{w} t$$

exercise: show there are no nontrivial ~~vector~~ constant vectors $\vec{w}$ that work here for all $t \in \mathbb{R}$.

So we try again: Create an idea of what the general solution should look like and see if we can find $\vec{w}$:

Let $\vec{x}^{(2)}(t) = \vec{v} t e^{-2t} + \vec{w} e^{-2t}$, and try to solve for both $\vec{v}$ and $\vec{w}$ at the same time.

We set again:

$$\frac{d}{dt} [\vec{v} t e^{-2t} + \vec{w} e^{-2t}] = A(\vec{v} t e^{-2t} + \vec{w} e^{-2t})$$

$$\vec{v} e^{-2t} - 2\vec{v} t e^{-2t} - 2\vec{w} e^{-2t} = A\vec{v} t e^{-2t} + A\vec{w} e^{-2t}$$

$$\vec{v} - 2\vec{w} - 2\vec{v} t = A\vec{v} t + A\vec{w}$$

These 2 sides are polynomials in t :

The coefficients in t : $-2\vec{v} = A\vec{v}$

This is solved precisely when $\vec{v}$ is an eigenvector: $\vec{v} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$.

The constant coefficients: ~~$A\vec{w}$~~ $\vec{v} - 2\vec{w} = A\vec{w}$

Rearrange to get $(A - (-2I))\vec{w} = \vec{v}$.

In this last equation, we call $\vec{w}$ a generalized eigenvector:

$$\left(\begin{bmatrix} -2 & 1 \\ 0 & -2 \end{bmatrix} - \begin{bmatrix} -2 & 0 \\ 0 & -2 \end{bmatrix} \right) \begin{bmatrix} w_1 \\ w_2 \end{bmatrix} = \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}$$

$$\Rightarrow \begin{cases} 0w_1 + 1w_2 = v_1 \\ 0w_1 + 0w_2 = v_2 \end{cases} \Rightarrow w_2 = 1, w_1 = \text{anything.}$$

choose $\vec{w} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. Then our guess

$$\vec{x}^{(2)}(t) = \vec{v}te^{-2t} + \vec{w}e^{-2t} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}te^{-2t} + \begin{bmatrix} 0 \\ 1 \end{bmatrix}e^{-2t}$$

is another solution to $\vec{x}' = \begin{bmatrix} -2 & 1 \\ 0 & -2 \end{bmatrix} \vec{x}$.

Q1: Is it a solution? Yes because we constructed it.

Q2: Is it independent of the other one?

$$\vec{x}^{(1)}(t) = \begin{bmatrix} e^{-2t} \\ 0 \end{bmatrix}, \quad \vec{x}^{(2)}(t) = \begin{bmatrix} te^{-2t} \\ e^{-2t} \end{bmatrix}$$

$$W(\vec{x}^{(1)}, \vec{x}^{(2)}) = \begin{vmatrix} e^{-2t} & te^{-2t} \\ 0 & e^{-2t} \end{vmatrix} = e^{-4t} \neq 0 \text{ on } \mathbb{R}.$$

Yes, independent.

Hence our general solution is

$$\vec{x}(t) = c_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} e^{-2t} + c_2 \left(\begin{bmatrix} 1 \\ 0 \end{bmatrix} t e^{-2t} + \begin{bmatrix} 0 \\ 1 \end{bmatrix} e^{-2t} \right)$$


To recap: Given $\vec{x}' = A\vec{x}$, where A has a repeated eigenvalue r and only 1 linearly indep. eigenvector $\vec{v}$, then the general solution is

$$\vec{x}(t) = c_1 \vec{v} e^{rt} + c_2 (\vec{v} t e^{rt} + \vec{w} e^{rt})$$

where $\vec{w}$ ~~solves~~ solves $(A - rI_2)\vec{w} = \vec{v}$.

What does this solution look like?

- Only straight line motion is along single eigenvector line.



- Other vector $\vec{w}$ helps to fill out the other dimension, but with the there is no motion along it that is straight
- Called a degenerate node, stable when $r < 0$ unstable when $r > 0$.
or an improper node.