

Spanning and Linear Independence

References are to Anton–Rorres, 7th edition

Coordinates Let V be a given vector space. We wish to equip V with a coordinate system, much as we did geometrically for the plane and space. We have the origin $\mathbf{0}$. However, because V is only a vector space, the concepts of length and orthogonality do not apply.

Take any set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors in V . There is an associated linear transformation $L: \mathbf{R}^r \rightarrow V$ (well hidden in Anton–Rorres), given by

$$L(k_1, k_2, \dots, k_r) = k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \dots + k_r\mathbf{v}_r \quad (1)$$

So $L(\mathbf{e}_i) = \mathbf{v}_i$ for each i . It is easy to check that L is linear. The idea is to choose S to make L an isomorphism of vector spaces, which will allow us to transfer everything from the general vector space V to the familiar vector space \mathbf{R}^r .

DEFINITION 2 The set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors in V is a *basis* [plural: *bases*] of V if the above linear transformation (1) satisfies the two conditions:

- (i) The *range* $R(L)$ of L is the whole of V ;
- (ii) The *kernel* $\text{Ker}(L)$ of L is $\{\mathbf{0}\}$.

Then by Theorem 8.3.1, L is 1–1 and we can restate the definition explicitly.

THEOREM 3 (=Thm. 5.4.1) *If $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ is a basis of V , then any vector $\mathbf{v} \in V$ can be uniquely expressed as a linear combination*

$$\mathbf{v} = k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \dots + k_r\mathbf{v}_r = L(k_1, k_2, \dots, k_r) \quad \square \quad (4)$$

We then have the *inverse* linear transformation $L^{-1}: V \rightarrow \mathbf{R}^r$ and $L^{-1}(\mathbf{v}_i) = \mathbf{e}_i$.

DEFINITION 5 If S is a basis of V , we define the *coordinate vector relative to S* of any vector $\mathbf{v} \in V$ to be $L^{-1}(\mathbf{v})$, and write it $(\mathbf{v})_S$. This is a vector in \mathbf{R}^r .

Explicitly, if \mathbf{v} is the linear combination (4), then $(\mathbf{v})_S = (k_1, k_2, \dots, k_r)$.

Example We have the standard basis $S = \{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_r\}$ of \mathbf{R}^r . In this case, L is the identity linear transformation and $(\mathbf{v})_S = \mathbf{v}$.

We break up Definition 2 and discuss the two conditions separately.

Spanning In any case, the range $R(L)$ of L is always a subspace of V .

DEFINITION 6 For any set S in V , we define the *span* of S to be the range $R(L)$ of the linear transformation L in equation (1), and write $\text{span}(S) = R(L)$.

Explicitly, $\text{span}(S)$ is the set of all linear combinations (4). Many different sets of vectors S can span the same subspace. Clearly, we can omit the zero vector $\mathbf{0}$ if it is present in S . More generally, as a direct application of Theorem 5.2.4, we have the following reduction, known as the Minus Theorem.

LEMMA 7 (=Thm. 5.4.4(b)) *Suppose $\mathbf{v}_i \in S$ is a linear combination of the other vectors in S . Let S' denote the set S with \mathbf{v}_i removed. Then $\text{span}(S') = \text{span}(S)$. \square*

Linear independence

DEFINITION 8 The set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ is *linearly independent* if the kernel $\text{Ker}(L)$ of the linear transformation L in equation (1) is $\{\mathbf{0}\}$, i.e. L is 1–1 (see Thm. 8.3.1). Otherwise, S is *linearly dependent*. [As linear independence is clearly the desirable condition, we shall eschew the term “linearly dependent”.]

Explicitly, S is linearly independent if there is no *linear relation*

$$k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \dots + k_r\mathbf{v}_r = \mathbf{0} \quad (9)$$

between the \mathbf{v} 's, other than the obvious *trivial* relation

$$0\mathbf{v}_1 + 0\mathbf{v}_2 + \dots + 0\mathbf{v}_r = \mathbf{0}$$

The following property is clear enough, but note the direction of the implication.

LEMMA 10 Let $T: V \rightarrow W$ be a linear transformation and $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ a set of vectors in V . If the image set $T(S) = \{T(\mathbf{v}_1), T(\mathbf{v}_2), \dots, T(\mathbf{v}_r)\}$ is linearly independent in W , then S is linearly independent in V .

Proof Suppose the vectors in S satisfy the linear relation

$$k_1\mathbf{v}_1 + k_2\mathbf{v}_2 + \dots + k_r\mathbf{v}_r = \mathbf{0}$$

We apply T to this to see that $T(S)$ satisfies the corresponding linear relation

$$k_1T(\mathbf{v}_1) + k_2T(\mathbf{v}_2) + \dots + k_rT(\mathbf{v}_r) = \mathbf{0} \quad \square$$

We need to recast the definition of linear independence in a more useful form. Roughly stated, S is linearly independent if each vector in S is new in the sense that it cannot be expressed in terms of the previous members of S .

LEMMA 11 (=Thm. 5.3.1(b), but sharper) The set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors is linearly independent if and only if $\mathbf{v}_1 \neq \mathbf{0}$ and no vector $\mathbf{v}_i \in S$ is a linear combination of the preceding vectors in S , i.e. for $2 \leq i \leq r$, $\mathbf{v}_i \notin \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{i-1}\}$.

Proof Assume there is a nontrivial linear relation (9), with k_m as the last nonzero coefficient. Then we can divide by k_m and rearrange equation (9) as

$$\mathbf{v}_m = l_1\mathbf{v}_1 + l_2\mathbf{v}_2 + \dots + l_{m-1}\mathbf{v}_{m-1} \quad (12)$$

where each $l_i = -k_i/k_m$. This expresses \mathbf{v}_m as a linear combination of the preceding vectors. (If $m = 1$, equation (12) degenerates to $\mathbf{v}_1 = \mathbf{0}$.) Conversely, if equation (12) holds for some m , we can rearrange it as the linear relation

$$l_1\mathbf{v}_1 + l_2\mathbf{v}_2 + \dots + l_{m-1}\mathbf{v}_{m-1} - \mathbf{v}_m = \mathbf{0} \quad \square$$

COROLLARY 13 (=Thm. 5.4.4(a), the Plus Theorem) Suppose the set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors in V is linearly independent but does not span V . Take any vector $\mathbf{v}_{r+1} \notin \text{span}(S)$. Then the enlarged set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r, \mathbf{v}_{r+1}\}$ remains linearly independent.

Proof The condition of Lemma 11 holds for \mathbf{v}_i if $i \leq r$ because S is linearly independent. It holds for \mathbf{v}_{r+1} by hypothesis. \square

Occasionally, a variant of Lemma 11 is useful.

COROLLARY 14 *The set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors in V is linearly independent if and only if $\mathbf{v}_r \neq \mathbf{0}$ and for $1 \leq i < r$, \mathbf{v}_i is not a linear combination of the later vectors in S .*

Proof We simply write the set S in reverse order and apply Lemma 11. \square

Vectors in \mathbf{R}^n All the main results depend ultimately on the following fact, which is intuitively obvious but *not* trivial to prove. However, the real work has already been done. Roughly, the consequence is that in a given vector space, a spanning set of vectors cannot be too small, and a linearly independent set cannot be too large.

LEMMA 15 (=Thm. 5.3.3) *A linearly independent set S of vectors in \mathbf{R}^n has at most n members.*

Proof Suppose S has r members, and consider the linear transformation $L: \mathbf{R}^r \rightarrow \mathbf{R}^n$ in equation (1). We are given $\text{Ker}(L) = \{\mathbf{0}\}$. Let A be the matrix of L , so that $L(\mathbf{x}) = A\mathbf{x}$. We know $A\mathbf{x} = \mathbf{0}$ has no nontrivial solutions. Since A is an $n \times r$ matrix, Theorem 1.2.1 shows that we must have $r \leq n$. \square

From this we deduce the result we really want.

THEOREM 16 *Suppose the vector space V is spanned by a set containing n vectors. Then any linearly independent set of vectors in V contains at most n members.*

Proof From the given spanning set, we construct as in equation (1) a linear transformation $L: \mathbf{R}^n \rightarrow V$ such that $R(L) = V$. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ be any linearly independent set of vectors in V . Since $R(L) = V$, we can choose for each i a vector $\mathbf{u}_i \in \mathbf{R}^n$ such that $L(\mathbf{u}_i) = \mathbf{v}_i$. Then by Lemma 10, the set $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_r\}$ is linearly independent in \mathbf{R}^n . Lemma 15 now shows that $r \leq n$. \square

COROLLARY 17 (=Thm. 5.4.3) *Any two bases of V contain the same number of vectors.* \square

DEFINITION 18 A vector space V is *finite-dimensional* if it has a basis that contains n vectors for some finite n . The number n is the *dimension* of V and is written $\dim(V)$. (By Corollary 17, it is well defined. For completeness, the zero vector space is considered to have dimension 0, and the *empty* set (not $\{\mathbf{0}\}$) as a basis; this works.)

We say that V is *infinite-dimensional* if it does not have a finite basis.

We are primarily interested in finite-dimensional vector spaces.

THEOREM 19 *Every finite-dimensional vector space is isomorphic to the standard vector space \mathbf{R}^n for a unique integer n .* \square

We collect in one place all the information about subsets of V .

THEOREM 20 (=Thms. 5.4.2 and 5.4.5) *Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ be any set of r vectors in the n -dimensional vector space V . Then:*

- (a) *If $r < n$, S does not span V . (It may or may not be linearly independent.)*
- (b) *If $r = n$, S spans V if and only if it is linearly independent. Thus S is a basis of V if either of these conditions holds.*
- (c) *If $r > n$, S is not linearly independent. (S may or may not span V .)*

Proof Parts (a) and (c) both follow immediately from Theorem 16.

For (b), suppose first that S spans V . If S is not linearly independent, Lemma 11 shows that some $\mathbf{v}_i \in S$ is a linear combination of the other members. We remove \mathbf{v}_i from S to get a set S' of $n-1$ vectors. By Lemma 7, S' still spans V ; but this contradicts (a).

Conversely, suppose that S is linearly independent. If S does not span V , we could use Corollary 13 to add another vector to S to form a linearly independent set of $n+1$ vectors in V , which would contradict (c). \square

It is not immediately obvious that any subspace of a finite-dimensional vector space is finite-dimensional.

THEOREM 21 (=Thm. 5.4.7) *If W is a subspace of the finite-dimensional vector space V , then W is again finite-dimensional and $\dim(W) \leq \dim(V)$, with equality only if $W = V$.*

Proof Choose a linearly independent set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors in W with r as large as possible; such a set exists because these vectors also lie in V , so that by Theorem 20(c), $r \leq n$ where $n = \dim(V)$. Then $\text{span}(S) = W$, otherwise Corollary 13 (applied in W) would allow us to extend S by one more vector and increase r by 1. So S must be a basis of W and $\dim(W) = r$.

If $W \neq V$, choose any vector $\mathbf{v}_{r+1} \in V$ that is not in W ; by Corollary 13, $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r, \mathbf{v}_{r+1}\}$ is still linearly independent. By Theorem 20(c), $r+1 \leq n$. \square

Constructing bases Obviously, if a subset S of V spans V , so does any subset S' of V that contains S . Any subset of a linearly independent set S remains linearly independent. Beyond these restrictions, we can construct bases of V as follows.

THEOREM 22 (=Thm. 5.4.6(a)) *Suppose the set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ spans the vector space V . Then we can thin out S to find a subset $S' \subset S$ that is a basis of V . In particular, V is finite-dimensional. (Explicitly, one possibility is to take S' as the set of all the nonzero $\mathbf{v}_i \in S$ that are not linear combinations of the preceding members of S , but there are other choices.)*

Proof If $\mathbf{v}_i \in S$ is a linear combination of the other members of S , we can delete \mathbf{v}_i from S without affecting $\text{span}(S)$, by Lemma 7. We repeat this, deleting elements of S one at a time, until we can go no further. By Lemma 11, the end result S' is linearly independent and therefore a basis of V . (We leave the suggested candidate for S' as an exercise.) \square

THEOREM 23 (=Thm. 5.4.6(b)) *Suppose given any linearly independent set $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ of vectors in a finite-dimensional vector space V that does not span V . Then we can extend S to a basis $S' = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r, \dots, \mathbf{v}_n\}$ of V .*

Proof By Corollary 13, we extend S by one more vector to get a larger subset that is still linearly independent. We repeat as long as possible, until we find a linearly independent set S' that does span V and is therefore a basis. The process must terminate, because by Theorem 20(c), the set S' can never have more than n members, where $n = \dim(V)$. (In fact, it has exactly n , by Corollary 17.) \square