

RIGIDITY OF NONNEGATIVELY CURVED SURFACES RELATIVE TO A CURVE

MOHAMMAD GHOMI AND JOEL SPRUCK

ABSTRACT. We prove that any properly oriented $\mathcal{C}^{2,1}$ isometric immersion of a positively curved Riemannian surface M into Euclidean 3-space is uniquely determined, up to a rigid motion, by its values on any curve segment in M . A generalization of this result to nonnegatively curved surfaces is presented as well under suitable conditions on their parabolic points. Thus we obtain a local version of Cohn-Vossen's rigidity theorem for convex surfaces subject to a Dirichlet condition. The proof employs in part Hormander's unique continuation principle for elliptic PDEs. Our approach also yields a short proof of Cohn-Vossen's theorem via Hopf's maximum principle.

1. INTRODUCTION

One of the fundamental results of classical surface theory is Cohn-Vossen's rigidity theorem [6, 7, 36, 38], which states that isometric closed nonnegatively curved surfaces in Euclidean 3-space are congruent. If the surface is not closed, however, it generally admits infinitely many noncongruent isometric immersions, and thus other constraints are needed to ensure its rigidity. Here we show that a local Dirichlet condition will suffice. For simplicity, we first state our main result for positively curved surfaces:

Theorem 1.1 (Main Theorem, First Version). *Let M be a connected 2-manifold and $f, \tilde{f}: M \rightarrow \mathbf{R}^3$ be $\mathcal{C}^{2,1}$ positively curved, isometric immersions whose mean curvature vectors induce the same orientation on M . Suppose that there exists a curve segment Γ in M and a proper rigid motion $\rho: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ such that $f = \rho \circ \tilde{f}$ on Γ . Then $f = \rho \circ \tilde{f}$ on M .*

In Section 5 below we will generalize the above theorem to nonnegatively curved surfaces, under suitable conditions on their parabolic points. The *manifold* M here may have boundary, and can assume any topological genus [15, 16]. *Isometric* means that the metrics induced on M by f and \tilde{f} coincide, i.e., $\langle df(v), df(w) \rangle = \langle d\tilde{f}(v), d\tilde{f}(w) \rangle$ for all tangent vectors $v, w \in T_p M$ and $p \in M$, where $\langle \cdot, \cdot \rangle$ is the standard inner product in \mathbf{R}^3 . If, furthermore, $df(v) \times df(w)$ is parallel to the mean

Date: May 15, 2018 (Last Typeset).

2010 Mathematics Subject Classification. Primary: 53A05, 35J96; Secondary 58J30, 52A15.

Key words and phrases. Isometric embedding, non-asymptotic curve, unique continuation, maximum principle, Darboux equation, Monge-Ampère equation, locally convex surface.

The research of M.G. was supported in part by NSF grant DMS-1711400.

curvature vector of f whenever $d\tilde{f}(v) \times d\tilde{f}(w)$ is parallel to the mean curvature vector of \tilde{f} , we say that the mean curvature vectors *induce the same orientation* on M . By a *curve segment* in M we mean the image of a smooth embedding $(-\epsilon, \epsilon) \rightarrow M$ (which may be arbitrarily small). Finally, a *proper rigid motion* is an orientation preserving isometry, i.e., $\rho \in \text{Iso}^+(\mathbf{R}^3) \simeq \mathbf{R}^3 \times \text{SO}(3)$.

The earliest antecedent to Theorem 1.1 appears to be a work of John Hewitt Jellett [28] who in 1849 studied how fixing a non-asymptotic curve in an analytic surface would render it infinitesimally rigid, see also Weingarten [40]. Later, in 1894, Darboux [9, Liv. 7, Chap. 5] established rigidity of analytic surfaces, relative to non-asymptotic curves, via Cauchy-Kovalevskaya theorem (see Notes 3.3 and 3.4). Indeed, non-asymptotic curves correspond to non-characteristic hypersurfaces for the underlying PDEs, and fixing a non-asymptotic curve in an isometric embedding fixes the derivatives of the embedding along that curve (see Note 3.2), which furnishes the Cauchy data. These notions are also implicit in the proofs of Cartan-Janet theorem [18, 26, 38] on analytic isometric embeddings.

As far as we know, Theorem 1.1 is the first analogue in the smooth category of the Jellett-Darboux rigidity result. Other results relevant to our work include a theorem of Alexandrov and Sen'kin [2], also see [33, p. 181], who showed that if a pair of isometric positively curved surfaces lie in the upper half-space, are star-shaped and concave with respect to the origin, and their corresponding boundary points are equidistant from the origin, then they are congruent. There is also a similar result of Pogorelov [33, p. 178] for convex caps which form concave graphs over the xy -plane, and whose corresponding boundary points have equal heights. For more background and references for rigidity problems in surface theory, which date back to Euler, Cauchy, and Maxwell, see [13, 18, 31, 33, 34, 38, 42].

The basic outline for proving Theorem 1.1 is as follows. After replacing \tilde{f} with $\rho \circ \tilde{f}$, we assume that $f = \tilde{f}$ on Γ and then aim to show that $f = \tilde{f}$ on M . To this end it suffices to establish that $f = \tilde{f}$ on an open neighborhood of a point of Γ (Section 2). This is achieved by showing first that f and \tilde{f} agree up to second order along Γ via geometric arguments (Section 3), and then applying a unique continuation principle for elliptic PDEs, with Lipschitz coefficients, due to Hormander (Section 4). Finally in Section 5 we will extend Theorem 1.1 to the nonnegative curvature case via works of Sacksteder [36] and Hartman-Nirenberg [19] on parabolic points of surfaces. These methods also yield a short proof of Cohn-Vossen's theorem, which is included in Appendix A.

Note 1.2 (Conditions of Theorem 1.1). The orientation condition in Theorem 1.1 is necessary. For instance let M be the upper hemisphere of \mathbf{S}^2 , f be the inclusion map, \tilde{f} be the reflection of f through the xy -plane, and Γ be any segment of the boundary of M . Further it is important that the curvature be positive, at least on Γ . Consider for instance a flat disk, and roll a corner of it outside its plane. According to [34, p.212, Rem. 7], there are even negatively curved isometric surfaces which coincide on an open set, but not everywhere else. Thus, in contrast to the Jellett-Darboux result, Theorem 1.1 appears to be a strictly elliptic phenomenon. Finally,

it is not necessary for f, \tilde{f} to be differentiable everywhere, but it is enough that they be continuous on M while they are $\mathcal{C}^{2,1}$ and isometric on $M \setminus X$, where X is any closed subset without interior points whose complement is connected and contains Γ . Then $f = \rho \circ \tilde{f}$ on $M \setminus X$ and therefore on M by continuity.

2. BEGINNING OF THE PROOF: LOCALIZATION

As mentioned above, we may replace \tilde{f} with $\rho \circ \tilde{f}$ so that $f = \tilde{f}$ on Γ . Furthermore, we may assume that Γ lies on the boundary ∂M of M . Indeed, if a point of Γ lies in the interior of the manifold, $\text{int}(M) := M \setminus \partial M$, we may extend a small segment containing that point to a closed curve $\bar{\Gamma}$ bounding a disk $D \subset \text{int}(M)$. Then $M' := M \setminus \text{int}(D)$ and D form a pair of manifolds whose boundaries contain Γ . So M will be rigid relative to Γ if and only if M' and D are rigid relative to Γ . To prove Theorem 1.1, it suffices then to show that:

Proposition 2.1. *Let M, f, \tilde{f} , and Γ be as in Theorem 1.1. Suppose that $\Gamma \subset \partial M$ and $\tilde{f} = f$ on Γ . Then every point of Γ has an open neighborhood in M where $f = \tilde{f}$.*

Indeed, suppose that the above proposition holds, let U be the union of all open sets in M where $f = \tilde{f}$, and ∂U be the topological boundary of U in M . Suppose, towards a contradiction, that there exists a point $q_1 \in \partial U \cap \text{int}(M)$, for otherwise we are done. Identify a neighborhood of q_1 in M with an open disk $\Omega \subset \mathbf{R}^2$ centered at q_1 . Let $q_0 \in U \cap \Omega$. There exists $\delta > 0$ such that the closed disk B_0 of radius δ centered at q_0 lies in $U \cap \Omega$. Let $q_t := (1-t)q_0 + tq_1$, and $s \in [0, 1]$ be the supremum of $t \in [0, 1]$ such that $B_t \subset U$, where B_t is the closed disk of radius δ centered at q_t . Then there exists a point $r \in \partial B_s \cap \partial U$. Applying Proposition 2.1 to a segment of ∂B_s containing r yields that $f = \tilde{f}$ on an open neighborhood of r . Thus $r \notin \partial U$, which is the desired contradiction. So it remains to prove Proposition 2.1, which is undertaken in the next two sections.

3. ORDER OF CONTACT ALONG Γ

We say that f and \tilde{f} have *contact of order 2* along Γ if, in some local coordinates, their derivatives agree up to second order on Γ . Here we show that, under the hypothesis of Proposition 2.1:

Lemma 3.1. *f and \tilde{f} have contact of order 2 along Γ .*

The above lemma appears to have been known, as a version of it is discussed in a Russian text by Kagan [29, p. 199–200]. We include our own treatment here, which will also yield a quick proof of the Jellett-Darboux theorem (Note 3.4).

To set the stage, we identify a small neighborhood Ω^+ of a point of Γ in M with a half disc in \mathbf{R}^2 bordering the y -axis, and lying to the right of it. We set $f(t) := f(0, t)$ for any mapping f defined on Ω^+ . Then, by assumption,

$$(1) \quad f(t) = \tilde{f}(t).$$

We need to show that the partial derivatives of f and \tilde{f} agree up to second order on Γ (the y -axis), i.e., $f_i(t) = \tilde{f}_i(t)$ and $f_{ij}(t) = \tilde{f}_{ij}(t)$ for $i, j = 1, 2$. To this end we first note that

$$f_2(t) = \tilde{f}_2(t), \quad \text{and} \quad f_{22}(t) = \tilde{f}_{22}(t).$$

By the isometry assumption we may also record that the coefficients of the induced metric tensor g of M are given by

$$(2) \quad g_{ij} := \langle f_i, f_j \rangle = \langle \tilde{f}_i, \tilde{f}_j \rangle =: \tilde{g}_{ij}.$$

Further we may assume that $\{f_1(t), f_2(t)\}$ and $\{\tilde{f}_1(t), \tilde{f}_2(t)\}$ are each orthonormal. This may be achieved by letting $\gamma(t)$ denote an arc length parametrization for Γ (with respect to g), $\nu(t)$ be the inward unit normal vector field along Γ (again with respect to g), and resetting

$$f(s, t) := f(\exp_{\gamma(t)}(s\nu(t))), \quad \text{and} \quad \tilde{f}(s, t) := \tilde{f}(\exp_{\gamma(t)}(s\nu(t))),$$

where \exp is the exponential map of M , and (s, t) ranges in a half-disk which we again denote by Ω^+ . Then $f(t)$ has unit speed, and $f_1(t), \tilde{f}_1(t)$ are inward conormals of $f(t)$ with respect to $f(\Omega^+), \tilde{f}(\Omega^+)$. More generally,

$$(3) \quad g_{11} = 1, \quad \text{and} \quad g_{12} = 0$$

on Ω^+ . These equations hold, via Gauss's Lemma, because $s \mapsto f(s, t)$ traces a geodesic with unit speed. Next note that since f has positive curvature, it has no asymptotic directions. So if

$$n(t) := f_1(t) \times f_2(t) \quad \text{and} \quad \tilde{n}(t) := \tilde{f}_1(t) \times \tilde{f}_2(t)$$

denote the unit normals of f and \tilde{f} on Γ , then $\langle f_{22}(t), n(t) \rangle$ and $\langle \tilde{f}_{22}(t), \tilde{n}(t) \rangle = \langle f_{22}(t), \tilde{n}(t) \rangle$ do not vanish. Further, by the orientation assumption in Theorem 1.1, they must have the same sign:

$$(4) \quad \langle f_{22}(t), n(t) \rangle \langle f_{22}(t), \tilde{n}(t) \rangle > 0.$$

Indeed, since the curvature is positive, the mean curvature vector points to the side of the tangent plane where the surface locally lies. Thus $\langle f_{22}(t), n(t) \rangle, \langle \tilde{f}_{22}(t), \tilde{n}(t) \rangle$ are both positive (negative) if and only if n, \tilde{n} are parallel (antiparallel) to the mean curvature vectors of f, \tilde{f} respectively.

3.1. First order contact. As we already know that $f_2(t) = \tilde{f}_2(t)$, it remains to check that $f_1(t) = \tilde{f}_1(t)$. Since $f(t)$ has unit speed, and $f_1(t), \tilde{f}_1(t)$ are inward conormals, the geodesic curvature of Γ with respect to the interior of M is given by

$$\langle f_{22}(t), f_1(t) \rangle = -\frac{1}{2}(g_{22})_1(t) = -\frac{1}{2}(\tilde{g}_{22})_1(t) = \langle \tilde{f}_{22}(t), \tilde{f}_1(t) \rangle.$$

By (4), $f_{22}(t) \neq 0$. So the principal normal $N(t) := f_{22}(t)/|f_{22}(t)|$ of $f(t)$ is well defined, and the last displayed expression yields that

$$(5) \quad \langle f_1, N \rangle = \langle \tilde{f}_1, N \rangle.$$

Now if $B(t) := f_2(t) \times N(t)$ denotes the binormal vector of $f(t)$, then $\{N(t), B(t)\}$ forms an orthonormal basis for the normal planes of $f(t)$, which contain $f_1(t)$. Thus

$$\langle f_1, N \rangle^2 + \langle f_1, B \rangle^2 = |f_1|^2 = |\tilde{f}_1|^2 = \langle \tilde{f}_1, N \rangle^2 + \langle \tilde{f}_1, B \rangle^2.$$

So it follows that $\langle f_1, B \rangle = \pm \langle \tilde{f}_1, B \rangle$. If $\langle f_1, B \rangle = -\langle \tilde{f}_1, B \rangle$, then

$$\langle n, N \rangle = \langle f_2 \times f_1, f_2 \times B \rangle = \langle f_1, B \rangle = -\langle \tilde{f}_1, B \rangle = -\langle f_2 \times \tilde{f}_1, f_2 \times B \rangle = -\langle \tilde{n}, N \rangle,$$

which contradicts (4). So we conclude that $\langle f_1, B \rangle = \langle \tilde{f}_1, B \rangle$ which together with (5) yields that

$$(6) \quad f_1(t) = \tilde{f}_1(t).$$

3.2. Second order contact. To show that the second derivatives of f and \tilde{f} match up along Γ first note that, since $g_{ij} = \tilde{g}_{ij}$,

$$\langle f_{ij}, f_k \rangle = \Gamma_{ij}^k = \frac{1}{2} \sum_{\ell} g^{\ell k} ((g_{\ell i})_j + (g_{j\ell})_i - (g_{ij})_{\ell}) = \tilde{\Gamma}_{ij}^k = \langle \tilde{f}_{ij}, f_k \rangle,$$

where $\Gamma_{ij}^k, \tilde{\Gamma}_{ij}^k$ are the Christoffel symbols associated to f, \tilde{f} , and $(g^{ij}) := (g_{ij})^{-1}$. So it remains to check that the coefficients of the second fundamental form $\ell_{ij} := \langle f_{ij}, n \rangle, \tilde{\ell}_{ij} := \langle \tilde{f}_{ij}, n \rangle$ agree on Γ . To this end note that

$$\ell_{12}(t) = -\langle f_1(t), n_2(t) \rangle = -\langle \tilde{f}_1(t), n_2(t) \rangle = \tilde{\ell}_{12}(t).$$

Further $\ell_{22}(t) = \tilde{\ell}_{22}(t)$, since $f_{22}(t) = \tilde{f}_{22}(t)$. By Theorema Egregium, the curvature $K := \det(\ell_{ij}) / \det(g_{ij})$ of f coincides with the curvature $\tilde{K} := \det(\tilde{\ell}_{ij}) / \det(\tilde{g}_{ij})$ of \tilde{f} . Indeed, Theorema Egregium does hold for \mathcal{C}^2 surfaces [20]. Thus

$$\det(\ell_{ij}) = K \det(g_{ij}) = \tilde{K} \det(\tilde{g}_{ij}) = \det(\tilde{\ell}_{ij}).$$

Furthermore, by (4), $\ell_{22}, \tilde{\ell}_{22}$ do not vanish along Γ . So

$$\ell_{11}(t) = \frac{\ell_{12}^2(t)}{\ell_{22}(t)} = \frac{\tilde{\ell}_{12}^2(t)}{\tilde{\ell}_{22}(t)} = \tilde{\ell}_{11}(t).$$

Hence $\ell_{ij}(t) = \tilde{\ell}_{ij}(t)$, as desired, which completes the proof of Lemma 3.1.

Note 3.2 (Non-asymptotic curves). In the proof of Lemma 3.1 above we used the positive curvature assumption only to ensure that (4) holds. Thus Lemma 3.1 holds for any pairs of \mathcal{C}^2 surfaces, regardless of their curvature, as long as Γ is *non-asymptotic*, i.e., never tangent to an asymptotic direction, and the normal curvatures of f, \tilde{f} assume the same sign along Γ .

Note 3.3 (The analytic case). By differentiating (2), and using (3), one quickly obtains the following equations, assuming that f is \mathcal{C}^3 , see Spivak [38, p. 150] or Han-Hong [18, p. 6]:

$$(7) \quad \langle f_{11}, f_2 \rangle = 0, \quad \langle f_{11}, f_1 \rangle = 0, \quad \langle f_{11}, f_{22} \rangle = -\frac{1}{2}(g_{22})_{11} + |f_{12}|^2.$$

By Cauchy-Kovalevskaya theorem, these equations have a unique solution once $f(t)$ and $f_1(t)$ have been prescribed, and f is analytic, see [38, p. 150–153] or [18, Lem. 1.1.4]. Thus, by (1) and (6), $f = \tilde{f}$ when \tilde{f} is also analytic. This proves Theorem 1.1 in the analytic case, and more generally establishes the rigidity of all analytic surfaces relative to non-asymptotic curves, as first observed by Darboux [9, p. 280]; see also Hopf and Samelson [22, Sec. 3]. For some isometric extension results in the smooth category see [14, 25, 27].

Note 3.4 (Higher order contact). When f and \tilde{f} are \mathcal{C}^k , it can be shown directly that they coincide up to order k along Γ . This yields a quick proof of Jellett-Darboux rigidity theorem for analytic surfaces, without invoking the Cauchy-Kovalevskaya theorem. Indeed, we claim that if f and \tilde{f} agree up to order $2 \leq m < k$ on Γ , then they agree up to order $m + 1$. To see this let $\alpha := \alpha_1 \alpha_2 \dots \alpha_m$, where $\alpha_i := 1, 2$. Then, $f_\alpha(t) = \tilde{f}_\alpha(t)$, for all α , which yields that $f_{\alpha 2}(t) = \tilde{f}_{\alpha 2}(t)$. By commutativity, it remains then to check that $f_{\alpha 1}(t) = \tilde{f}_{\alpha 1}(t)$, where all $\alpha_i = 1$. Since $\{f_1(t), f_2(t), f_{22}(t)\}$ is linearly independent, due to the non-asymptotic assumption on Γ , this follows from repeatedly differentiating the equations (7) with respect to the first variable, which shows that $f_{\alpha 1}(t)$ is determined by $f_{\alpha' 2}(t)$, $f_\alpha(t)$, and lower order derivatives. Thus, by induction, f and \tilde{f} agree up to order k on Γ . Consequently, $f = \tilde{f}$ on M when f and \tilde{f} are analytic.

4. UNIQUE CONTINUATION

To complete the proof of Proposition 2.1, and therefore of Theorem 1.1, it remains to show that $f = \tilde{f}$ on the region Ω^+ discussed in the last section. To this end, let Ω^- be the reflection of Ω^+ with respect to the y -axis, and set $\Omega := \Omega^+ \cup \Omega^-$. We may extend f, \tilde{f} isometrically to all of Ω , without losing regularity, as follows. By the Lipschitz version of Whitney's extension theorem [4, Thm. 2.64], first we extend f to Ω so that $f \in \mathcal{C}^{2,1}(\Omega, \mathbf{R}^3)$; see also [11, p. 10] or [10, Sec. 5.4] for explicit constructions via “higher order reflection”. Then we extend \tilde{f} by setting it equal to f on Ω^- . By Lemma 3.1, f agrees with \tilde{f} up to order 2 on the y -axis, so \tilde{f}_{ij} are continuous on Ω ; furthermore, \tilde{f}_{ij} are Lipschitz both on Ω^+ and Ω^- , which quickly yields that \tilde{f}_{ij} are Lipschitz on Ω . So $\tilde{f} \in \mathcal{C}^{2,1}(\Omega, \mathbf{R}^3)$ as well.

After replacing Ω by a smaller disc, we may assume that f and \tilde{f} are positively curved on Ω . Now for a unit vector $e \in \mathbf{R}^3$, let $u := \langle f, e \rangle$, $\tilde{u} := \langle \tilde{f}, e \rangle$. Then $u, \tilde{u} \in \mathcal{C}^{2,1}(\Omega)$, and they both satisfy the Darboux equation [18, p. 45]:

$$(8) \quad \det(\nabla_{ij} u) = K \det(g_{ij})(1 - |\nabla u|_g^2),$$

where (∇_{ij}) and ∇u are the Riemannian Hessian and gradient respectively and $|\cdot|_g := \sqrt{\langle \cdot, \cdot \rangle_g}$ is the Riemannian norm. More explicitly,

$$\nabla_{ij} u := u_{ij} - \sum_k \Gamma_{ij}^k u_k, \quad \text{and} \quad \nabla u := \sum_{ij} g^{ij} u_i \partial_j,$$

where ∂_j denote the standard basis of \mathbf{R}^2 . Thus (8) is a fully nonlinear Monge-Ampère equation of general form [39, Sec. 3.8], and is elliptic since $K > 0$ [18, p. 46]. We claim that $\phi := u - \tilde{u}$ vanishes identically on Ω , which is all we need. Indeed, since e was chosen arbitrarily, $u \equiv \tilde{u}$ implies that $f \equiv \tilde{f}$. To this end, we subtract (8) from its counterpart in terms of \tilde{u} . A straight forward computation yields that

$$(9) \quad \sum_{ij} (\nabla_{ij}^* u + \nabla_{ij}^* \tilde{u}) \nabla_{ij} \phi + 2K \det(g_{ij}) \langle \nabla(u + \tilde{u}), \nabla \phi \rangle_g = 0,$$

where $(\nabla_{ij}^*) := \det(\nabla_{ij})(\nabla_{ij})^{-1}$ is the cofactor matrix of (∇_{ij}) , i.e., $\nabla_{11}^* = \nabla_{22}$, $\nabla_{12}^* = -\nabla_{21}$, and $\nabla_{22}^* = \nabla_{11}$. Note that (9) is linear in terms of ϕ , and ϕ vanishes on an open subset of Ω . Hence, by the following unique continuation principle, see also Armstrong and Silvestre [3, Prop. 2.1] and Garofolo and Lin [12], ϕ vanishes identically on Ω as claimed.

Lemma 4.1 (Hormander [24], Thm. 17.2.6). *Let $\Omega \subset \mathbf{R}^2$ be a connected domain and $u: \Omega \rightarrow \mathbf{R}$ be a solution to the linear equation*

$$(10) \quad \sum_{ij} a_{ij}(x) u_{ij} + \sum_i b_i(x) u_i + c(x) u = 0,$$

where $a_{ij}: \Omega \rightarrow \mathbf{R}$ are uniformly elliptic and Lipschitz, while $b_i, c: \Omega \rightarrow \mathbf{R}$ are bounded measurable functions. If u vanishes on an open subset of Ω , then $u \equiv 0$.

Uniformly elliptic means that the eigenvalues of (a_{ij}) are bounded below by a positive constant. To check this and other requirements needed to apply Lemma 4.1 to (9) note that in this context

$$a_{ij} = \nabla_{ij}^* u + \nabla_{ij}^* \tilde{u}.$$

So a_{ij} are Lipschitz, since $u, \tilde{u} \in \mathcal{C}^{2,1}(\Omega)$. Next note that $\det(\nabla_{ij} u), \det(\nabla_{ij} \tilde{u}) > 0$ by (8), since $K > 0$ by assumption. Thus eigenvalues of $(\nabla_{ij} u)$ and $(\nabla_{ij} \tilde{u})$ never vanish on Ω . Now since, by construction, u, \tilde{u} agree up to second order at some point of Ω , these eigenvalues will coincide at one point, and thus will always carry the same sign. In particular we may assume that they are positive on Ω , after replacing e with $-e$ if necessary. So $(\nabla_{ij} u)$ and $(\nabla_{ij} \tilde{u})$ are positive definite matrices. Consequently their cofactor matrices $(\nabla_{ij}^* u)$ and $(\nabla_{ij}^* \tilde{u})$ are positive definite as well. Hence so is their sum (a_{ij}) . Now we may assume that a_{ij} are uniformly elliptic on Ω , after replacing Ω by a smaller disk with compact closure $\overline{\Omega}' \subset \Omega$. Finally note that b_i are continuous on $\overline{\Omega}'$, while $c \equiv 0$, so they are all bounded and measurable. This concludes the proof of Theorem 1.1.

5. NONNEGATIVE CURVATURE

Here we generalize Theorem 1.1 to nonnegatively curved surfaces. Note that if the set of parabolic, or zero curvature, points $M^0 \subset M$ of a nonnegatively curved $\mathcal{C}^{2,1}$ immersion $f: M \rightarrow \mathbf{R}^3$ does not have interior points and does not disconnect M , then $f = \tilde{f}$ on $M \setminus M^0$ by Theorem 1.1, and therefore, by continuity, $f = \tilde{f}$ on M . Thus the nontrivial case is when $\text{int}(M^0)$ is nonempty.

Theorem 5.1 (Main Theorem, Full Version). *Let M , f and \tilde{f} be as in Theorem 1.1, except that the curvature of f is allowed to be nonnegative. Let $M^0 \subset M$ be the set of points where the curvature of f vanishes. Suppose that*

- (i) M^0 contains no curve ℓ such that $f(\ell)$ is a complete line,
- (ii) M^0 is complete, i.e., its Cauchy sequences converge,
- (iii) $M^0 \subset \text{int}(M)$,
- (iv) $M \setminus M^0$ is connected.

If there exists a curve segment Γ in $M \setminus M^0$ and a proper rigid motion $\rho: \mathbf{R}^3 \rightarrow \mathbf{R}^3$ such that $f = \rho \circ \tilde{f}$ on Γ , then $f = \rho \circ \tilde{f}$ on M .

To prove this result, we will again replace \tilde{f} with $\rho \circ \tilde{f}$, so that $f = \tilde{f}$ on Γ , and show that $f = \tilde{f}$ on M . Let $M^+ := M \setminus M^0$, and $\overline{M^+}$ be the closure of M^+ in M . Since M^+ is connected, Theorem 1.1 yields that $f = \tilde{f}$ on M^+ , and therefore on $\overline{M^+}$ by continuity. In particular it follows that the second fundamental forms of f and \tilde{f} agree on $\overline{M^+}$. Further since $M^0 \subset \text{int}(M)$, we have $\partial M^0 \subset \overline{M^+}$. Now the following result, which requires conditions (i) and (ii) above, immediately completes the proof of Theorem of 5.1, via the fundamental theorem of surfaces.

Lemma 5.2 (Sacksteder [36], Thm. I). *If the second fundamental forms of f and \tilde{f} agree on ∂M^0 , then they agree on M^0 .*

Since Sacksteder's argument is somewhat involved, we include here a short simple proof of Theorem 5.1 for the case where M^0 is compact (in which case conditions (i) and (ii) are automatically satisfied). Recall that we just need to check that $f = \tilde{f}$ on $\text{int}(M^0)$. To this end, let $M^F \subset \text{int}(M^0)$ be the set of points with a *flat neighborhood*, i.e., a neighborhood mapped by f into a plane. The following fact from Spivak [37] is implicit in the works of Hartman-Nirenberg [19] and Massey [30] on \mathcal{C}^2 surfaces of zero curvature. See also Pogorelov [33, p. 609] or [35, p. 79] for an extension of this fact to the \mathcal{C}^1 category.

Lemma 5.3 ([37], Cor. 8, p. 243). *Through every point $p \in \text{int}(M^0) \setminus M^F$ there passes a curve ℓ with end point(s) on ∂M^0 such that f maps ℓ homeomorphically into a straight line segment or ray in \mathbf{R}^3 .*

When M^0 is compact, ℓ has finite length, and so it has two end points q_i . Further, since $q_i \in \partial M^0 \subset \overline{M^+}$, $f(q_i) = \tilde{f}(q_i)$. So $\tilde{f}(\ell)$ is a curve joining $f(q_i)$, with the same arc length as $f(\ell)$ by isometry. Consequently $f = \tilde{f}$ on ℓ , which yields $f = \tilde{f}$ on $\text{int}(M^0) \setminus M^F$. Next let C be a component of M^F . Then $\partial C \subset \overline{M^+} \cup (M^0 \setminus M^F)$. So $f = \tilde{f}$ on ∂C , which yields that $f = \tilde{f}$ on C . Indeed, through each point $p \in C$ there passes a curve ℓ with end points $q_i \in \partial C$ such that $f(\ell)$ is a line segment (let L be a complete line passing through $f(p)$ in the plane of $f(C)$, and ℓ be the closure of the component of $f^{-1}(L) \cap C$ containing p). Hence, again $f = \tilde{f}$ on ℓ , since $f(q_i) = \tilde{f}(q_i)$. So $f = \tilde{f}$ on M^F , and consequently on M^0 .

Note 5.4 (The case of zero curvature). The above argument shows that if M is compact, and $f, \tilde{f}: M \rightarrow \mathbf{R}^3$ are \mathcal{C}^2 isometric immersions with everywhere vanishing

curvature, then $f = \tilde{f}$ on M whenever $f = \tilde{f}$ on ∂M . In other words, compact developable surfaces in \mathbf{R}^3 are rigid relative to their boundary.

APPENDIX A. A SHORT PROOF OF COHN-VOSSEN'S THEOREM

Cohn-Vossen proved the first version of his rigidity result in 1927 for positively curved analytic surfaces [6], before extending it to \mathcal{C}^3 surfaces in 1936 [7], see Hopf [23, p. 168]. The proof of this theorem included in various texts, e.g., [5, 18, 38], is the 1943 argument by Herglotz [21] based on his celebrated integral formula. Later Wintner [41] established the theorem for \mathcal{C}^2 surfaces, and Sacksteder [36] extended it to nonnegative curvature; see also [8, Sec. 6.3] and [17]. Following the same outline as in the proof of Theorem 1.1, we present a proof of Cohn-Vossen's theorem which is even shorter than Herglotz's and works immediately in the \mathcal{C}^2 category.

Let $f, \tilde{f}: \mathbf{S}^2 \rightarrow \mathbf{R}^3$ be \mathcal{C}^2 positively curved isometric immersions, with principal curvatures $k_1 \leq k_2, \tilde{k}_1 \leq \tilde{k}_2$ respectively. By the invariance of Gauss curvature, $k_1 k_2 \equiv \tilde{k}_1 \tilde{k}_2$. So if, at some point, $k_2 < \tilde{k}_2$, then $\tilde{k}_1 < k_1$, which in turn yields that $\tilde{k}_1 < \tilde{k}_2$. As is well-known, it is impossible for the last inequality to hold everywhere; because then the principal directions of \tilde{f} corresponding to \tilde{k}_2 , would generate a line field on \mathbf{S}^2 , in violation of the Poincaré-Hopf index theorem [37, Thm. 20, p. 223]. So we conclude that $k_2(p) = \tilde{k}_2(p)$ for some point $p \in \mathbf{S}^2$, which in turn yields that $k_1(p) = \tilde{k}_1(p)$. Consequently, after a rigid motion, we may assume that f and \tilde{f} have contact of order 2 at p .

Let $\Omega \subset \mathbf{R}^2$ be an open disk, and $\theta: \overline{\Omega} \rightarrow \mathbf{S}^2$ be a smooth map with $\theta(\partial\Omega) = p$ such that $\theta: \Omega \rightarrow \mathbf{S}^2 \setminus \{p\}$ is a diffeomorphism. Replace f, \tilde{f} by $f \circ \theta, \tilde{f} \circ \theta$ respectively. Further, as in Section 4, set $u := \langle f, e \rangle, \tilde{u} := \langle \tilde{f}, e \rangle$ for a unit vector $e \in \mathbf{R}^3$. Then $u, \tilde{u} \in \mathcal{C}^2(\overline{\Omega})$, and $u = \tilde{u}$ on $\partial\Omega$. Again, as in Section 4, set $\phi := u - \tilde{u}$. Then ϕ satisfies the linear elliptic equation (9), which can be put in the form (10) with $c \equiv 0$. Now, since $\phi = 0$ on $\partial\Omega$, Hopf's maximum principle [10, p. 332] yields that $\phi = 0$ on Ω . So $u \equiv \tilde{u}$ and, since e was arbitrary, we conclude that $f \equiv \tilde{f}$, as desired.

The above argument might also work for nonnegative curvature via an appropriate version of the maximum principle for degenerate equations. In closing, we should recall that Pogorelov [32, 33] generalized Cohn-Vossen's theorem to all closed convex surfaces regardless of their regularity in 1952, although that proof remains long and intricate. See also Volkov [1, Sec. 12.1] for another approach to Pogorelov's theorem via uniform rigidity estimates.

ACKNOWLEDGMENTS

We thank Igor Belegradek, Robert Bryant, Jeanne Clelland, Robert Greene, Idjad Sabitov, Andrzej Swiech, and Deane Yang for useful communications. Jellett's work [28] and its citation in [29] were first pointed out to us by Sabitov. Further we are grateful to Sabitov for suggesting a simplification of our original proof of Theorem 5.1 in the compact case.

REFERENCES

- [1] A. D. Alexandrov, *Convex polyhedra*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2005. Translated from the 1950 Russian edition by N. S. Dairbekov, S. S. Kutateladze and A. B. Sossinsky, With comments and bibliography by V. A. Zalgaller and appendices by L. A. Shor and Yu. A. Volkov. MR2127379 (2005j:52002) ↑9
- [2] A. D. Alexandrov and E. P. Sen'kin, *On the rigidity of convex surfaces*, Vestnik Leningrad. Univ. **10** (1955), no. 8, 3–13. MR0072500 ↑2
- [3] S. N. Armstrong and L. Silvestre, *Unique continuation for fully nonlinear elliptic equations*, Math. Res. Lett. **18** (2011), no. 5, 921–926. MR2875864 ↑7
- [4] A. Brudnyi and Y. Brudnyi, *Methods of geometric analysis in extension and trace problems. Volume 1*, Monographs in Mathematics, vol. 102, Birkhäuser/Springer Basel AG, Basel, 2012. MR2882877 ↑6
- [5] S. S. Chern, *Curves and surfaces in Euclidean space*, Studies in global geometry and analysis, 1967, pp. 16–56. MR35#3610 ↑9
- [6] S. E. Cohn-Vossen, *Zwei sätze über die starrheit der eiflächen*, Nach. Gesellschaft Wiss. Göttinger, Math Phys. Kl. (1927), 125–134. ↑1, 9
- [7] ———, *Bending of surfaces in the large*, Uspekhi Mat. Nauk **1** (1936), 33–76. ↑1, 9
- [8] M. Dajczer, *Submanifolds and isometric immersions*, Mathematics Lecture Series, vol. 13, Publish or Perish, Inc., Houston, TX, 1990. Based on the notes prepared by Mauricio Antonucci, Gilvan Oliveira, Paulo Lima-Filho and Rui Tojeiro. MR1075013 ↑9
- [9] G. Darboux, *Leçons sur la théorie générale des surfaces et les applications géométriques du calcul infinitésimal. Troisième partie*, Chelsea Publishing Co., Bronx, N. Y., 1972. Lignes géodésiques et courbure géodésique. Paramètres différentiels. Déformation des surfaces, Réimpression de la première édition de 1894. MR0396213 ↑2, 6
- [10] L. C. Evans, *Partial differential equations*, Graduate Studies in Mathematics, vol. 19, American Mathematical Society, Providence, RI, 1998. MR1625845 ↑6, 9
- [11] A. Friedman, *Partial differential equations*, Holt, Rinehart and Winston, Inc., New York-Montreal, Que.-London, 1969. MR0445088 ↑6
- [12] N. Garofalo and F.-H. Lin, *Unique continuation for elliptic operators: a geometric-variational approach*, Comm. Pure Appl. Math. **40** (1987), no. 3, 347–366. MR882069 ↑7
- [13] M. Ghomi, *Open problems in geometry of curves and surfaces*, www.math.gatech.edu/~ghomi/Papers/op.pdf. ↑2
- [14] M. Ghomi and R. E. Greene, *Relative isometric embeddings of Riemannian manifolds*, Trans. Amer. Math. Soc. **363** (2011), no. 1, 63–73. MR2719671 (2011k:53072) ↑6
- [15] M. Ghomi and M. Kossowski, *h-principles for hypersurfaces with prescribed principal curvatures and directions*, Trans. Amer. Math. Soc. **358** (2006), no. 10, 4379–4393 (electronic). MR2231382 ↑1
- [16] H. Gluck and L.-H. Pan, *Embedding and knotting of positive curvature surfaces in 3-space*, Topology **37** (1998), no. 4, 851–873. MR98m:57008 ↑1
- [17] P. Guan and X. S. Shen, *A rigidity theorem for hypersurfaces in higher dimensional space forms*, Analysis, complex geometry, and mathematical physics: in honor of Duong H. Phong, 2015, pp. 61–65. MR3372460 ↑9
- [18] Q. Han and J.-X. Hong, *Isometric embedding of Riemannian manifolds in Euclidean spaces*, Mathematical Surveys and Monographs, vol. 130, American Mathematical Society, Providence, RI, 2006. MR2261749 (2008e:53055) ↑2, 5, 6, 7, 9
- [19] P. Hartman and L. Nirenberg, *On spherical image maps whose Jacobians do not change sign*, Amer. J. Math. **81** (1959), 901–920. MR0126812 ↑2, 8
- [20] P. Hartman and A. Wintner, *On the fundamental equations of differential geometry*, Amer. J. Math. **72** (1950), 757–774. MR0038110 ↑5
- [21] G. Herglotz, *über die Starrheit der Eiflächen*, Abh. Math. Sem. Hansischen Univ. **15** (1943), 127–129. MR0014714 ↑9

- [22] H. Hopf and H. Samelson, *Zum Beweis des Kongruenzsatzes für Eiflächen*, Math. Z. **43** (1938), no. 1, 749–766. MR1545746 ↑6
- [23] H. Hopf, *Differential geometry in the large*, Second, Springer-Verlag, Berlin, 1989. Notes taken by Peter Lax and John W. Gray, With a preface by S. S. Chern, With a preface by K. Voss. MR90f:53001 ↑9
- [24] L. Hörmander, *The analysis of linear partial differential operators. III*, Classics in Mathematics, Springer, Berlin, 2007. Pseudo-differential operators, Reprint of the 1994 edition. MR2304165 ↑7
- [25] N. Hungerbühler and M. Wasem, *The one-sided isometric extension problem*, Results Math. **71** (2017), no. 3-4, 749–781. MR3648443 ↑6
- [26] T. A. Ivey and J. M. Landsberg, *Cartan for beginners*, Graduate Studies in Mathematics, vol. 175, American Mathematical Society, Providence, RI, 2016. MR3586335 ↑2
- [27] H. Jacobowitz, *Extending isometric embeddings*, J. Differential Geometry **9** (1974), 291–307. MR0377773 ↑6
- [28] J. H. Jellett, *On the properties of inextensible surfaces*, Trans. Irish. Ac. Dublin **22** (1849), 343–377. ↑2, 9
- [29] V. F. Kagan, *Osnovy teorii poverhnostei v tenzornom izloženii. čast' vtoraya. Poverhnosti v prostranstve. Oobraženiya i izgibaniya poverhnostei. Special' nye voprosy (Russian)*, Gosudarstv. Izdat. Tehn.-Teor. Lit., Moscow-Leningrad, 1948. MR0049606 ↑3, 9
- [30] W. S. Massey, *Surfaces of Gaussian curvature zero in Euclidean 3-space*, Tôhoku Math. J. (2) **14** (1962), 73–79. MR0139088 ↑8
- [31] I. Pak, *Lectures on discrete and polyhedral geometry*, www.math.ucla.edu (2008). ↑2
- [32] A. V. Pogorelov, *Odnosnačnaya opredelennost' obščih vypuklyh poverhnostei [unique determination of general convex surfaces.]*, Monografii Instituta Matematiki, vyp. II, Akad. Nauk Ukrainської SSR, Kiev, 1952. MR0063681 ↑9
- [33] ———, *Extrinsic geometry of convex surfaces*, American Mathematical Society, Providence, R.I., 1973. MR0346714 (49 #11439) ↑2, 8, 9
- [34] I. Kh. Sabitov, *Local theory of bendings of surfaces*, Geometry, III, 1992, pp. 179–256. MR1306736 ↑2
- [35] ———, *Isometric immersions and embeddings of locally Euclidean metrics*, Reviews in Mathematics and Mathematical Physics, vol. 13, Cambridge Scientific Publishers, Cambridge, 2008. MR2584444 ↑8
- [36] R. Sacksteder, *The rigidity of hypersurfaces*, J. Math. Mech. **11** (1962), 929–939. MR0144286 ↑1, 2, 8, 9
- [37] M. Spivak, *A comprehensive introduction to differential geometry. Vol. III*, Second, Publish or Perish Inc., Wilmington, Del., 1979. MR82g:53003c ↑8, 9
- [38] ———, *A comprehensive introduction to differential geometry. Vol. V*, Second, Publish or Perish Inc., Wilmington, Del., 1979. MR532834 (82g:53003e) ↑1, 2, 5, 6, 9
- [39] N. S. Trudinger and X.-J. Wang, *The Monge-Ampère equation and its geometric applications*, Handbook of geometric analysis. No. 1, 2008, pp. 467–524. MR2483373 ↑7
- [40] J. Weingarten, *Ueber die Deformationen einer biegsamen unausdehnbaren Fläche*, J. Reine Angew. Math. **100** (1887), 296–310. MR1580097 ↑2
- [41] A. Wintner, *On isometric surfaces*, Amer. J. Math. **74** (1952), 198–214. MR0047351 ↑9
- [42] S.-T. Yau, *Review of geometry and analysis*, Mathematics: frontiers and perspectives, 2000, pp. 353–401. MR1754787 (2001m:53003) ↑2

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GA 30332

E-mail address: ghomi@math.gatech.edu

URL: www.math.gatech.edu/~ghomi

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD 21218

E-mail address: js@math.jhu.edu

URL: www.math.jhu.edu/~js