

## Math 107, Fall 2006: Midterm II Practice Exam 1 Solutions

### Solutions

1. This question concerns the function

$$f(x, y) = 1 - yx^2.$$

- (a) Draw the cross-section through the graph of  $f(x, y)$  given by setting  $y = 1$ .
- (b) Calculate the partial derivative of  $f$  with respect to  $x$  and evaluate it at the point  $(2, 1)$ .
- (c) On your graph from (a), draw a tangent line to the cross-section whose slope is represented by your answer from (b).
- (a) The graph should be of  $z$  against  $x$  with  $z = 1 - x^2$  (from setting  $y = 1$ ).
- (b)  $\frac{\partial f}{\partial x} = -2xy$ , which at the point  $(2, 1)$  is  $-4$ .
- (c) Part (b) tells us that the slope of the graph at  $x = 2$  should be  $-4$ .
2. (a) Draw a graph showing the  $c$ -level curves of the function  $f(x, y) = x^2 - y$  for  $c = -2, 0, 2$ .
- (b) Mark on your graph from (a) the direction of the gradient vector  $\nabla f$  at the points  $(0, 0)$ ,  $(2, 6)$  and  $(-1, -1)$ . (Note: you do not have to get the right length for these gradient vectors. It is the direction that is important.)
- (c) What is the directional derivative of  $f(x, y)$  in the direction  $(1, 0)$  at the point  $(0, 0)$ ? (You must give a reason or a calculation.)
- (a) The  $c$ -level curve is the curve with equation  $x^2 - y = c$ , or  $y = x^2 - c$ . So the level curves are parabolas.
- (b) The gradient vectors should be perpendicular to the level curves, and should point in the direction of increasing  $f$ .
- (c) The directional derivative can be calculated as

$$D_{\mathbf{u}}f(0, 0) = \frac{\partial f}{\partial x}(0, 0)u_1 + \frac{\partial f}{\partial y}(0, 0)u_2.$$

The partial derivatives are:

$$\frac{\partial f}{\partial x} = 2x, \quad \frac{\partial f}{\partial y} = -1$$

So the directional derivative is:

$$0 \times 1 + (-1) \times 0 = 0.$$

Alternatively, you could just say that the vector  $(1, 0)$  is perpendicular to the gradient vector at  $(0, 0)$ , and so the directional derivative in that direction is zero.

3. Find the critical point of the function  $f(x, y) = x^2 + 3y^2$ . Is this a local max, local min or saddle? (Show your work.)

The critical point occurs where both partial derivatives are zero. So, we need  $2x = 0$  and  $6y = 0$ . Therefore,  $x = 0$  and  $y = 0$ . So the critical point is  $(0, 0)$ .

To see if this is a local max/min or saddle, we calculate the Hessian matrix:

$$Hf = \begin{pmatrix} 2 & 0 \\ 0 & 6 \end{pmatrix}.$$

This has determinant  $+12$ , so  $(0, 0)$  is either a local max or min. We then check the top-left and bottom-right entries. These are both positive, so the critical point is a local minimum.

4. Calculate each of the following partial derivatives:

(a) if  $f(x, y) = e^{x^2y}$ , find  $\frac{\partial^2 f}{\partial y^2}$ ;

(b) if  $f(x, y) = \sin(2x + y)$ , find  $\frac{\partial^2 f}{\partial x \partial y}$ ;

(a)  $\frac{\partial f}{\partial y} = x^2 e^{x^2y}$  and so  $\frac{\partial^2 f}{\partial y^2} = x^4 e^{x^2y}$ .

(b)  $\frac{\partial f}{\partial y} = \cos(2x + y)$  and so  $\frac{\partial^2 f}{\partial x \partial y} = -2 \sin(2x + y)$ .

5. Use the chain rule to find  $f'(t)$  where  $f(x, y) = x^2y$ ,  $x(t) = \sin t$ ,  $y(t) = e^t$ . (Your answer should be in terms of  $t$  only.)

The chain rule says that

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

In our case:  $\partial f/\partial x = 2xy$ ,  $dx/dt = \cos t$ ,  $\partial f/\partial y = x^2$  and  $dy/dt = e^t$ . Therefore

$$f'(t) = 2xy \cos t + x^2 e^t.$$

Substituting in the expressions for  $x$  and  $y$ , we get:

$$f'(t) = 2e^t \sin t \cos t + e^t (\sin t)^2.$$