Name .....

## PRACTICE EXAM 2 40pts.

- There are 6 pages in the exam including this page.
- Write all your answers clearly. You have to show work to get points for your answers.
- You can write on both sides of the paper. Indicate that the answer follows on the back of the page.
- Use of Calculators is *not* allowed during the exam.
- $(1) \dots /8$
- $(2) \dots /8$
- $(3) \dots /8$
- $(4) \ldots / 16$

Total  $\ldots ... /40$ 

(1) *8pts.* Let  $T: I\!\!R^4 \to I\!\!R^4$  be defined by a matrix

$$A = \begin{bmatrix} 1 & 0 & -2 & 2 \\ 0 & 1 & -1 & 0 \\ 3 & 2 & 0 & 1 \\ 1 & -1 & 0 & 4 \end{bmatrix}$$

(a) Find the Kernel of T.

(b) Is T invertible? Why or why not?

(2a) *3pts* When is a set of vectors  $\{v_1, \cdots, v_n\}$  in a vector space V said to be a basis of V?

(2b) 5pts. Let 
$$A = \begin{bmatrix} 1 & 2 & 0 & 1 \\ 0 & 4 & 1 & 2 \\ -1 & 2 & 1 & 1 \\ 2 & 0 & -1 & 0 \end{bmatrix}$$
 represent a linear transformation from  $T$ :  
 $I\!\!R^4 \to I\!\!R^4$ . Find a basis for the Im $T$ .

(3) *8pts.* Check if the following sets are subspaces of  $\mathbb{R}^3$  and  $\mathbb{R}^4$  respectively or not. Explain your answers.

(a) 
$$W = \left\{ \begin{bmatrix} x \\ y \\ z \end{bmatrix} : \begin{array}{c} x - y + z = 0 \\ x + 1 - 2z = 0 \end{array} \right\}$$

(b) 
$$V = \left\{ \begin{bmatrix} 0\\ a+b\\ c\\ c-5a \end{bmatrix} : a, b, c \in \mathbb{R} \right\}.$$

- (4) 16pts. Give short answers to the following.
  - (a) If A and B are  $2 \times 2$  matrices such that AB = 0 then, either A = 0 or B = 0.

(b) Suppose that A is a  $3 \times 3$  matrix such that Ax = x for all  $x \in \mathbb{R}^3$ . Let  $I_3$  be the  $3 \times 3$  identity matrix. Find  $\text{Ker}(A - I_3)$ , that is the Kernel of the transformation represented by  $A - I_3$ . (c) Let  $T : \mathbb{R}^2 \to \mathbb{R}^2$  be the transformation which maps a vector  $v \in \mathbb{R}^2$  to its reflection along a line along  $\begin{bmatrix} 1\\1 \end{bmatrix}$ . Describe the matrix of this transformation.

(d) Is 
$$\left\{ \begin{bmatrix} 1\\3\\-1 \end{bmatrix}, \begin{bmatrix} 3\\2\\0 \end{bmatrix}, \begin{bmatrix} 4\\5\\-1 \end{bmatrix} \right\}$$
 a basis of  $\mathbb{R}^3$ ? Why or why not?