

GLOBAL EXISTENCE OF FREE-ENERGY SOLUTIONS TO THE 2D PATLAK–KELLER–SEGEL–NAVIER–STOKES SYSTEM WITH CRITICAL AND SUBCRITICAL MASS

CHEN-CHIH LAI, JUNCHENG WEI, AND YIFU ZHOU

ABSTRACT. We consider a coupled Patlak–Keller–Segel–Navier–Stokes system in \mathbb{R}^2 that describes the collective motion of cells and fluid flow, where the cells are attracted by a chemical substance and transported by ambient fluid velocity, and the fluid flow is forced by the friction induced by the cells. The main result of the paper is to show the global existence of free-energy solutions to the 2D Patlak–Keller–Segel–Navier–Stokes system with critical and subcritical mass.

1. INTRODUCTION

In this paper, we consider the Cauchy problem of the following 2D Patlak–Keller–Segel–Navier–Stokes system (PKS–NS system) [43]

$$\begin{cases} \partial_t n + u \cdot \nabla n = \Delta n - \nabla \cdot (n \nabla c) & \text{in } \mathbb{R}^2 \times (0, \infty), \\ -\Delta c = n & \text{in } \mathbb{R}^2 \times (0, \infty), \\ \partial_t u + u \cdot \nabla u + \nabla P = \Delta u + n \nabla c, \quad \nabla \cdot u = 0 & \text{in } \mathbb{R}^2 \times (0, \infty), \\ (n, u)(\cdot, 0) = (n_0, u_0) & \text{in } \mathbb{R}^2, \end{cases} \quad (1.1)$$

where n , c , u and P represent the cell density, the concentration of chemoattractant, fluid velocity field and pressure, respectively, and (n_0, u_0) is a given initial data such that $\nabla \cdot u_0 = 0$. The system (1.1)₁–(1.1)₂ is the parabolic-elliptic Patlak–Keller–Segel equations with additional effect of advection by a shear flow [4, 5, 44], which models the chemotaxis phenomena in a moving fluid. The equations (1.1)₃ is known as the forced Navier–Stokes equations describing the fluid motion subject to forcing generated by the cells. The forcing $n \nabla c$, which also appears in the Nernst–Planck–Navier–Stokes system [28] with an opposite sign, is to make the cells move without acceleration and to match the aggregation nonlinearity in the cell density evolution system.

The system (1.1) can be viewed as a coupling between the Patlak–Keller–Segel system (PKS system) and the incompressible Navier–Stokes equations (NS equations). Many significant contributions have been made on the studies to the Patlak–Keller–Segel system and the incompressible Navier–Stokes equations. The Patlak–Keller–Segel system is first introduced by Patlak [72] and Keller and Segel [54] and simplified by Nanjundiah [70]. We refer to [49, 50] for a review of the mathematical problems in chemotaxis models. It is well known that the PKS system is L^1 critical [51] and the L^1 norm of the solution is conserved. When the initial measure has a small atomic part, a unique mild solution is constructed in [8]. There is a competition between the tendency of cells to spread all over \mathbb{R}^2 by diffusion and the tendency to aggregate due to the drift caused by the chemoattractivity. The balance between these two mechanisms happens exactly at the critical mass $M = 8\pi$. In fact, in the subcritical case $M < 8\pi$, the

global well-posedness of the free energy solution of PKS system is established in [19, 32] under the assumptions of finite initial second moment and finite initial entropy. The proof relies on the free energy functional introduced in [10, 34, 69] and the logarithmic Hardy–Littlewood–Sobolev inequality. We refer the reader to [24] for an alternative proof using Delort’s theory on the 2D incompressible Euler equation [30]. For the asymptotic behavior of the solutions in the subcritical case, we refer the interested reader to [18, 22]. On the other hand, for the case of supercritical mass $M > 8\pi$, under suitable assumption, the solution blow up in finite time due to the evolution equation of second moment, see [9–11, 13, 19, 26, 33, 45–47, 64, 65, 74, 81] for example. Let us mention that the continuation of measure-valued solutions of the PKS system in the sense of Poupaud’s weak solutions [73], which make sense even if there are mass concentrations, have also been studied (see [6, 14, 33, 61, 82, 84]). As for the critical case $M = 8\pi$, solutions are shown to be global in time (see [12] for the radial case and [83] for the general case). An infinite time blow-up free energy solution having a finite second moment was found in [17]. The solution converges to a Dirac measure as time goes to infinity. If $M \leq 8\pi$ and the entropy of the initial data is finite, the global-in-time existence of classical solutions of finite entropy without the assumption of finite second moment is proved in [68] (also see [31] in the radial case). A classical survey can be found in [15]. The precise description of the infinite-time dynamic and the limiting profile is given in [25, 39, 60]. Recently, it is shown in [29] that there exists a radial function n_0^* with mass 8π such that for any initial data n_0 sufficiently close to n_0^* the solution starting from n_0 is globally defined and blows up at infinite time, with precise blow-up behaviors and stability obtained. It is shown in [85] using a monotonicity formula that mild solutions of PKS exists globally in time if and only if $M \leq 8\pi$ without any additional assumptions. We refer readers to [16] for non-blow-up free energy solutions which have an infinite second moment. For the incompressible Navier–Stokes equation, the existence of global weak solutions to the initial value problem has been established by Leray [58] and Hopf [48]. For comprehensive results regarding the Navier–Stokes equation, we refer the interested readers to [20, 35, 57, 59, 75, 76, 78, 80], the references therein, and [1, 7, 21, 36–38, 40–42, 52, 53, 71] for results in two and general dimensions.

Similar to the PKS system, the PKS–NS system (1.1) formally preserves mass, in the sense that

$$\int_{\mathbb{R}^2} n(x, t) dx = \int_{\mathbb{R}^2} n_0(x) dx =: M$$

for all $t \in (0, \infty)$ because u is divergence-free. Moreover, since the solution of the two dimensional Poisson equation in (1.1)₂ is given up to a harmonic function, we directly define the concentration of the chemoattractant by

$$c(x, t) = \frac{1}{2\pi} \int_{\mathbb{R}^2} \log \frac{1}{|x - y|} n(x, t) dy.$$

Furthermore, the PKS–NS system (1.1) possesses a decreasing free energy functional [43, (1.2)]

$$\mathcal{F}[n, u] := \int_{\mathbb{R}^2} \left[n \left(\log n - \frac{1}{2} c \right) + \frac{1}{2} |u|^2 \right] dx, \quad (1.2)$$

where the first and the second term are the entropy and a potential energy of the density n , respectively, and the third term is the kinetic energy of the velocity field u . As shown in [43, Lemma 1.1], the free energy is dissipative along the dynamics (1.1).

Lemma 1.1 (Free Energy Functional). *Let (n, u) be a smooth solution of (1.1) and $\mathcal{F}[n, u]$ be given in (1.2), then*

$$\frac{d}{dt} \mathcal{F}[n, u] = - \int_{\mathbb{R}^2} [n |\nabla(\log n - c)|^2 + |\nabla u|^2] dx.$$

Proof. A direct computation shows

$$\frac{d}{dt} \mathcal{F}[n, u] = - \int_{\mathbb{R}^2} n |\nabla(\log n - c)|^2 dx + \int_{\mathbb{R}^2} nu \cdot \nabla c dx - \int_{\mathbb{R}^2} |\nabla u|^2 dx + \int_{\mathbb{R}^2} n \nabla c \cdot u dx.$$

The lemma follows by integrating the last term by parts and using $\nabla \cdot u = 0$. \square

In [43], the local and global existence of solutions to the Patlak–Keller–Segel–Navier–Stokes system (1.1) are established in the Sobolev space H^s , $s \geq 2$, when the initial mass is strictly less than 8π . The H^s bound of their solutions grows exponentially in time. They also consider (1.1) on a torus \mathbb{T}^2 , and prove a similar existence result as for \mathbb{R}^2 while the solution can be bounded in H^s uniformly in time. There are many open problems regarding the PKS–NS system (1.1). For example, it is unclear that solutions exists globally in time or there exists a finite-time singularity when $M \geq 8\pi$. The ambient fluid flow might suppress the potential blow-up in (1.1)₁–(1.1)₂ (see [4, 5, 44, 55]). Despite of the known results of suppression of explosion on torus, finite-time blow-ups of (1.1) are possible in \mathbb{R}^2 when u is divergence-free. In a forthcoming work [56], we construct infinite time blow-up solutions to the PKS–NS system in the critical case $M = 8\pi$ by a new inner-outer gluing method. It is worth mentioning that in the radially symmetric class, the PKS–NS system is decoupled which can be seen from the dynamics of the second moment (1.7) and PKS system itself can develop finite time singularities in the case $M > 8\pi$ (see [27, 63, 81] for example). We will get back to the construction of finite time blow-up in the supercritical case $M > 8\pi$ for the PKS–NS system without any symmetry in a future work. In another aspect, the PKS–NS system in a bounded domain with Neumann boundary condition is also an intriguing problem.

The main aim of this paper is to show the existence of global-in-time solutions for the Patlak–Keller–Segel–Navier–Stokes system (1.1) with mass $M \leq 8\pi$. Throughout the paper, the initial data are assumed to satisfy

$$0 \leq n_0 \in L^1(\mathbb{R}^2), \quad n_0 \log n_0 \in L^1(\mathbb{R}^2), \quad n_0 \log(1 + |x|) \in L^1(\mathbb{R}^2), \quad (1.3)$$

and

$$u_0 \in W_{0,\sigma}^{1,2}(\mathbb{R}^2) := \overline{C_{0,\sigma}^\infty(\mathbb{R}^2)}^{\|\cdot\|_{W^{1,2}(\mathbb{R}^2)}} = \{u \in W^{1,2}(\mathbb{R}^2) : \operatorname{div} u = 0\}, \quad (1.4)$$

where $C_{0,\sigma}^\infty(\mathbb{R}^2) := \{u \in C^\infty(\mathbb{R}^2) : \operatorname{div} u = 0\}$.

We now define the free-energy solution of the PKS–NS system (1.1) of which the concept is introduced in [17] for PKS system.

Definition 1.2 (Free-Energy Solution). *Given $T > 0$, (n, u) is a free-energy solution to the Patlak–Keller–Segel–Navier–Stokes system (1.1) with initial data (n_0, u_0) on $[0, T]$ if $(1 + |\log n|)n \in L^\infty(0, T; L^1(\mathbb{R}^2))$, $u \in L^\infty(0, T; L^2(\mathbb{R}^2)) \cap L^2(0, T; \dot{H}^1(\mathbb{R}^2))$, (n, u) satisfies (1.1) in distributional sense, and*

$$\mathcal{F}[n, u](t) + \int_0^t \int_{\mathbb{R}^2} \left[n(x, s) |\nabla(\log n(x, s) - c(x, s))|^2 + \frac{1}{2} |\nabla u(x, s)|^2 \right] dx ds \leq \mathcal{F}[n_0, u_0] \quad (1.5)$$

for almost every $t \in (0, T)$.

It will be shown in Section 3 that every mild solution (n, u) is a free-energy solution.

The following is our main result on global-in-time existence of solutions to (1.1).

Theorem 1.3 (Global existence). *Suppose that (n_0, u_0) satisfies the assumptions (1.3) and (1.4), and*

$$M := \int_{\mathbb{R}^2} n_0 dx \leq 8\pi,$$

then the free-energy solution (n, u) to the Patlak–Keller–Segel–Navier–Stokes system (1.1) exists globally in time. Moreover, the L^∞ -norm of the solution does not blow up in finite time, i.e., for any $0 < t_0 < T < \infty$

$$\sup_{t \in (t_0, T)} (\|n(t)\|_\infty + \|u(t)\|_\infty) < \infty. \quad (1.6)$$

In order to prove Theorem 1.3, we need to use the following proposition.

Proposition 1.4. *Let (n_0, u_0) satisfy (1.3) and (1.4). Assume also $\|n_0\|_1 = 8\pi$. For a local-in-time mild solution (n, u) on $[0, T)$ with initial data (n_0, u_0) given in Theorem 3.2, it holds that for $0 < t_0 < T$,*

$$\sup_{t_0 \leq t < T} \int_{\mathbb{R}^2} (1 + n(t)) \log(1 + n(t)) dx < \infty.$$

Although the PKS–NS system shares many features with the PKS system because of u being divergence-free, there are still some significant properties holds for PKS but fails for PKS–NS. It is well-known that solutions of the PKS system with suitable spatial decay also satisfy the conservation of the first moment (the center of mass) $\int_{\mathbb{R}^2} n(x, t) x dx$, and the equation of the second moment (the variance) $\frac{d}{dt} \int_{\mathbb{R}^2} |x|^2 n(x, t) dx = 4M \left(1 - \frac{M}{8\pi}\right)$. From the evolution equation of the second moment, we can easily see that solutions with finite initial second moment do not exist globally if $M > 8\pi$. Moreover, the second moment is conserved for the critical case $M = 8\pi$, which is one of the keys to prove the global existence in [17]. However, for the PKS–NS system (1.1), unlike the PKS system, the dynamics of the first moment and the second moment are more delicate due to the coupling from the velocity field. In fact, they satisfy

$$\frac{d}{dt} \int_{\mathbb{R}^2} n(x, t) x_i dx = \int_{\mathbb{R}^2} n(x, t) u_i(x, t) dx$$

and

$$\frac{d}{dt} \int_{\mathbb{R}^2} |x|^2 n(x, t) dx = 4M - \frac{M^2}{2\pi} + 2 \int_{\mathbb{R}^2} n(x, t) u(x, t) \cdot x dx. \quad (1.7)$$

The sign of the last term in (1.7) is unknown. It is unclear how to control the integral unless the solution is radially symmetric [43, Corollary 1], in which case the integral vanishes (the system gets decoupled). The lack of control over the second moment is the main obstacle to bound solutions uniform in time. Instead of using the second moment, we employ the following modified free energy functional

$$H[n, u](t) := \int_{\mathbb{R}^2} (1 + n(t)) \log(1 + n(t)) dx - \frac{1}{2} \int_{\mathbb{R}^2} n(t) c(t) dx + \frac{1}{2} \int_{\mathbb{R}^2} |u(t)|^2 dx \quad (1.8)$$

and the Brezis–Merle inequality as in [68] to estimate the modified free energy in the interior region in the proof of Proposition 1.4. Details of the proof of Proposition 1.4 are given in Section 4. It is worth mentioning that the modified entropy $\int_{\mathbb{R}^2} (1 + n(t)) \log(1 + n(t)) dx$ in (1.8) is non-negative and has been used to get nonnegative global solutions for parabolic-parabolic Keller–Segel system in [67] and [62].

Remark 1.1. Unlike the free energy functional $\mathcal{F}[n, u]$, the modified energy $H[n, u]$ defined in (1.8) does not decrease in time. In fact, it formally satisfies

$$\frac{d}{dt} H[n, u] = - \int_{\mathbb{R}^2} \left| \nabla \log(1+n) - \frac{\nabla c}{2} \right|^2 dx - \int_{\mathbb{R}^2} n |\nabla \log(1+n) - \nabla c|^2 dx + \int_{\mathbb{R}^2} \frac{|\nabla c|^2}{2} dx.$$

Let us explain the idea for the proof of our main result, Theorem 1.3. First of all, we employ the free energy functional $\mathcal{F}[n, u]$ defined in (1.2) and the logarithmic Hardy–Littlewood–Sobolev inequality to obtain the a priori estimate of the fluid velocity. For the critical case $M = 8\pi$, the energy of fluid velocity is uniformly bounded in time (see Lemma 3.3), and thus the velocity of mild solution verify the Prodi–Serrin conditions for regularity (see Remark 3.3 and Remark 3.4). The regularity of u allows us to control all additional terms arising from the coupling when deriving the a priori entropy estimate of the advection–Patlak–Keller–Segel system (1.1)₁–(1.1)₂ in Section 4. For the subcritical regime $M < 8\pi$, we improve the result of [43] by reducing the regularity assumption on the initial data (n_0, u_0) . In fact, we show that solutions with the initial conditions (1.3) and (1.4) become H^s in short time, thus the global-in-time existence follows directly from [43].

The paper is organized as follows. In Section 2, we prepare some analysis tools that used in this paper. In Section 3, we introduce the notion of mild solutions of (1.1) and discuss local existence and regularity of the mild solutions. In Section 4, we derive the a priori estimate of the modified entropy and prove Proposition 1.4. Section 5 is devoted to prove Theorem 1.3, the global existence of solutions to (1.1).

2. PRELIMINARY

Before proving our main theorem, we recall some well-known and useful lemmas.

In the light of the critical Sobolev embedding in \mathbb{R}^2 , we define the class of functions of the bounded mean oscillations $BMO(\mathbb{R}^2)$ by

$$\|f\|_{BMO} := \sup_B \frac{1}{|B|} \int_B |f - f_B| dx \quad \text{with } f_B := \frac{1}{|B|} \int_B f dx,$$

where the supremum being taken over the set of Euclidean balls B and $|B|$ is the Lebesgue measure of B . It is well-known that $H^1(\mathbb{R}^2)$ is embedded in $BMO(\mathbb{R}^2) \cap L^2(\mathbb{R}^2)$ as a corollary of the Poincaré inequality.

In view of the modified entropy $\int_{\mathbb{R}^2} (1+n) \log(1+n) dx$ appears in the modified free energy, we recall the notion of the Orlicz space.

Definition 2.1 (Orlicz space [2, 79]). *Let $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a convex function such that*

$$\lim_{s \rightarrow 0_+} \frac{\phi(s)}{s} = 0, \quad \lim_{s \rightarrow \infty} \frac{\phi(s)}{s} = \infty. \quad (2.1)$$

Then the Orlicz class $L_\phi(\mathbb{R}^2)$ consists of all measurable functions $f : \mathbb{R}^2 \rightarrow \mathbb{C}$ satisfying

$$\int_{\mathbb{R}^2} \phi(|f(x)|) dx < \infty.$$

Since $\phi(s) = (1+s) \log(1+s) - s$ is a non-negative, convex function satisfying (2.1), non-negative functions with finite total mass and finite modified entropies forms the Orlicz class $L_{(1+\cdot) \log(1+\cdot)} =: L \log L$. If $f \in L \log L$, then the solution ψ to the Poisson equation

$$-\Delta \psi = f \quad \text{in } \mathbb{R}^2 \quad (2.2)$$

is a locally bounded function. If $f \log(1 + |x|) \in L^1$ then ψ becomes a locally integrable function.

The following proposition of *BMO* estimate for the solution of two dimensional Poisson equation is well known.

Proposition 2.2 (*BMO estimate*). *Let $f \log(2 + |x|) \in L^1(\mathbb{R}^2)$ and ψ be the solution of the Poisson equation (2.2). Then we have*

$$\|\psi\|_{BMO} \leq C \|f\|_1,$$

where C is a constant independent of f .

Proof. It is well known that $\log|x| \in BMO$. Since

$$\psi(x) - \frac{1}{|B|} \int_B \psi(y) dy = \int_{\mathbb{R}^2} \left[\log|x-y| - \frac{1}{|B|} \int_B \log|z-y| dz \right] f(y) dy,$$

we have

$$\begin{aligned} & \frac{1}{|B|} \int_B \left| \psi(x) - \frac{1}{|B|} \int_B \psi(y) dy \right| dx \\ & \leq \frac{1}{|B|} \int_{\mathbb{R}^2} \int_B \left| \log|x-y| - \frac{1}{|B|} \int_B \log|z-y| dz \right| dx |f(y)| dy \\ & = \frac{1}{|B|} \int_{\mathbb{R}^2} \int_{B_y} \left| \log|x| - \frac{1}{|B_y|} \int_{B_y} \log|z| dz \right| dx |f(y)| dy, \quad B_y = B + y, \\ & \leq \frac{1}{|B|} \|\log|\cdot|\|_{BMO} \|f\|_1, \end{aligned}$$

completing the proof. \square

BMO functions are locally L^p . In fact, using the John–Nirenberg inequality, one can prove the following lemma.

Lemma 2.3 ([77, (7), pp. 144]). *Suppose $f \in BMO(\mathbb{R}^2)$. Then for any $1 < p < \infty$, f is locally in L^p , and*

$$\frac{1}{|B|} \int_B |f - f_B|^p dx \leq C_p \|f\|_{BMO}^p$$

for any balls B .

We also make use of the logarithmic Hardy–Littlewood–Sobolev inequality to derive the energy estimate of the velocity field u .

Proposition 2.4 (Logarithmic Hardy–Littlewood–Sobolev Inequality [3, 23]). *Let f be a nonnegative function in $L^1(\mathbb{R}^2)$ such that $f \log f$ and $f \log(1 + |x|^2)$ belong to $L^1(\mathbb{R}^2)$. If $\int_{\mathbb{R}^2} f dx = M$, then*

$$\int_{\mathbb{R}^2} f \log f dx + \frac{2}{M} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} f(x) f(y) \log|x-y| dx dy \geq -C(M)$$

with $C(M) := M(1 + \log \pi - \log M)$.

The logarithmic Hardy–Littlewood–Sobolev inequality has been broadly applied to the PKS system. It is the key ingredient to show the global-in-time existence in [19] and to characterize the blowup profile in [17]. To be more precise, it is used to derive an upper bound of the entropy from the dissipative free energy functional for the PKS system.

As mentioned in the introduction, we employ the modified free energy functional $H[n, u]$ given in (1.8). The following lemma implies the equivalence of the free energy function $\mathcal{F}[n, u]$ and the modified one $H[n, u]$ under the weaker condition that $n_0 \log(2 + |x|) \in L^1$.

Lemma 2.5 ([68, Lemma 2.3]). *Let f be a nonnegative measurable function on a measurable set Ω in \mathbb{R}^n satisfying $f \log(2 + |x|) \in L^1(\Omega)$. Then*

$$f \log f \in L^1(\Omega) \quad \Leftrightarrow \quad (1 + f) \log(1 + f) \in L^1(\Omega).$$

In fact,

$$\int_{\Omega} (1 + f) \log(1 + f) dx \leq 2 \int_{\Omega} f |\log f| dx + (2 \log 2) \int_{\Omega} f dx$$

and

$$\int_{\Omega} f |\log f| dx \leq \int_{\Omega} (1 + f) \log(1 + f) dx + 2\alpha \int_{\Omega} f \log(2 + |x|) dx + \frac{1}{e} \int_{\Omega} \frac{1}{(2 + |x|)^\alpha} dx,$$

where $n < \alpha < \infty$.

The following lemma is the key to reach the global existence Theorem 1.3 from Proposition 1.4. It gives us a control of the $L_{t,x}^3$ norm of n in the energy equality of the density n .

Lemma 2.6 ([67, Lemma 2.1 (2)]). *For any $\varepsilon > 0$, there exists $C_\varepsilon > 0$ such that for $f \geq 0$*

$$\|f\|_2 \leq \varepsilon \|(1 + f) \log(1 + f)\|_1^{\frac{1}{2}} \|\nabla f\|_2^{\frac{1}{2}} + C(\varepsilon) \|f\|_1^{\frac{1}{2}} \quad (2.3)$$

$$\|f\|_3 \leq \varepsilon \|(1 + f) \log(1 + f)\|_1^{\frac{1}{3}} \|\nabla f\|_2^{\frac{2}{3}} + C(\varepsilon) \|f\|_1^{\frac{1}{3}} \quad (2.4)$$

where $C(\varepsilon) = O((e^{\varepsilon^{-3}} - 1)^{\frac{2}{3}})$ as $\varepsilon \rightarrow 0$.

In order to control the free energy in the interior region, we need the following lemma concerning the Poisson equation in \mathbb{R}^2 , which is a consequence of the Brezis–Merle inequality under the zero Dirichlet boundary condition.

Lemma 2.7 ([68, Lemma 2.7]). *Let Ω be a bounded domain in \mathbb{R}^2 with smooth boundary. For $g \in L^2(\Omega)$, let $v \in W^{2,2}(\Omega)$ be a solution of $-\Delta v = g$ in Ω . If $\|g\|_{L^1(\Omega)} < 4\pi$, then*

$$\int_{\Omega} \exp(|v(x)|) dx \leq \frac{4\pi^2}{4\pi - \|g\|_{L^1(\Omega)}} d(\Omega)^2 \exp\left(\sup_{\partial\Omega} |v(x)|\right)$$

where $d(\Omega)$ is the diameter of Ω .

It follows from the following lemma that the L^∞ -norm of ∇c is controlled by the L^q -norm of n , $q \in (2, \infty)$.

Lemma 2.8 ([66, Lemma 2.5]). *Let $2 < q \leq \infty$. For $f \in L^1 \cap L^q$,*

$$\left\| \nabla \frac{1}{2\pi} \log \frac{1}{|\cdot|} * f \right\|_{\infty} \leq C_q \|f\|_1^{\frac{q-2}{2(q-1)}} \|f\|_q^{\frac{q}{2(q-1)}},$$

where

$$C_q = (2\pi)^{1/2} \left(\frac{q-1}{q-2} \right)^{1/2} \left\{ \left(\frac{q}{q-1} \right)^{\frac{q-2}{2(q-1)}} + \left(\frac{q}{q-1} \right)^{-\frac{q}{2(q-1)}} \right\}.$$

3. LOCAL-IN-TIME EXISTENCE OF MILD SOLUTIONS

We begin with the definition of the mild solution of the PKS–NS system (1.1).

Definition 3.1 (Mild Solution). *Given $(n_0, u_0) \in L^1 \times L^2_{\sigma}$, we define $(n(t), u(t))$ to be a mild solution of the Patlak–Keller–Segel–Navier–Stokes system (1.1) on $[0, T]$ with initial data (n_0, u_0) if*

- (i) $n \in C([0, T]; L^1) \cap C((0, T); L^{4/3})$ and $u \in C([0, T]; L^2) \cap C((0, T); L^4)$,
- (ii) $\sup_{0 < t < T} t^{1/4} (\|n(t)\|_{4/3} + \|u(t)\|_4) < \infty$,
- (iii) (n, u) satisfies the integral equations

$$n(t) = e^{t\Delta} n_0 - \int_0^t e^{(t-s)\Delta} \nabla \cdot (n(s) \nabla c(s) + n(s) u(s)) ds, \quad (3.1)$$

$$u(t) = e^{t\Delta} u_0 - \int_0^t e^{(t-s)\Delta} \mathbf{P} \nabla \cdot (\nabla c(s) \otimes \nabla c(s) + u(s) \otimes u(s)) ds, \quad (3.2)$$

where $-\Delta c(s) = n(s)$.

In this section, we discuss the local existence and regularity of mild solutions to (1.1).

Let us write (3.1)–(3.2) as

$$(n(t), u(t)) = (e^{t\Delta} n_0, e^{t\Delta} u_0) - B((n, u), (n, u)),$$

where $B = (B_1, B_2)$ is a bilinear form in which B_1 and B_2 are bilinear forms defined by

$$B_1((m, v), (n, u)) := \int_0^t e^{(t-s)\Delta} \nabla \cdot (m(s) \nabla c(s) + m(s) u(s)) ds,$$

$$B_2((m, v), (n, u)) := \int_0^t e^{(t-s)\Delta} \mathbf{P} \nabla \cdot (\nabla b(s) \otimes \nabla c(s) + v(s) \otimes u(s)) ds$$

in which $-\Delta c(s) = n(s)$ and $-\Delta b(s) = m(s)$.

By the Hardy–Littlewood–Sobolev inequality, for any $q \in (2, \infty)$

$$\|\nabla c\|_q \lesssim \|n\|_{\frac{2q}{2+q}}. \quad (3.3)$$

We recall the classical L^q – L^p estimates of heat and Stokes semigroups. For any $1 \leq q \leq p \leq \infty$ ($p \neq 1, q \neq \infty$), there holds

$$\begin{aligned} \|e^{t\Delta} f\|_p + \|e^{t\Delta} \mathbf{P} f\|_p &\lesssim t^{\frac{1}{p} - \frac{1}{q}} \|f\|_q, \\ \|e^{t\Delta} \nabla \cdot F\|_p + \|e^{t\Delta} \mathbf{P} \nabla \cdot F\|_p &\lesssim t^{-\frac{1}{2} + \frac{1}{p} - \frac{1}{q}} \|F\|_q. \end{aligned} \quad (3.4)$$

Let \mathcal{E}_T be a Banach space defined as

$$\mathcal{E}_T = \left\{ (n, u) \in L^\infty(0, T; L^1) \times L^\infty(0, T; L^2) : \right. \\ \left. t^{1/4}(n(\cdot, t), u(\cdot, t)) \in L^\infty(0, T; L^{4/3}) \times L^\infty(0, T; L^4) \right\}$$

with the norm

$$\|(n, u)\|_{\mathcal{E}_T} := \sup_{t \in (0, T)} t^{1/4} \left(\|n(\cdot, t)\|_{4/3} + \|u(\cdot, t)\|_4 \right) + \sup_{t \in (0, T)} (\|n(\cdot, t)\|_1 + \|u(\cdot, t)\|_2) \\ := \|(n, u)\|_{X_T} + \|(n, u)\|_{Y_T}.$$

We first establish the local-in-time existence of mild solutions to (1.1) and some important properties of the solutions in the following theorem.

Theorem 3.2. *Given $n_0 \in L^1$ and $u_0 \in L^2_\sigma$, there exists $T \in (0, \infty)$ such that the Cauchy problem (1.1) has a unique mild solution (n, u) on $[0, T)$. Moreover, (n, u) satisfies the following properties:*

- (i) $n(t) \rightarrow n_0$ in L^1 and $u(t) \rightarrow u_0$ in L^2 ;
- (ii) for every $1 \leq p \leq \infty$, there holds $n \in C((0, T]; L^p)$ and $\sup_{0 < t < T} (t^{1-1/p} \|n(t)\|_p) < \infty$;
- (iii) for every $1 \leq p \leq 2$, there holds $u \in C((0, T]; L^p)$ and $\sup_{0 < t < T} (t^{1-1/p} \|u(t)\|_{\frac{2p}{2-p}}) < \infty$;
- (iv) for every $m \in \mathbb{Z}_+$, $l \in \mathbb{Z}_+^2$ and $1 < p < \infty$, there holds $\partial_t^m \partial_x^l n, \partial_t^m \partial_x^l u \in C((0, T]; L^p)$;
- (v) (n, u) is a classical solution of (1.1) in $\mathbb{R}^2 \times (0, T)$;
- (vi) If $n_0 \geq 0$ and $n_0 \neq 0$, then $n(x, t) > 0$ for all $(x, t) \in \mathbb{R}^2 \times (0, T)$;
- (vii) If $n_0 \log(1 + |x|) \in L^1$, then $n(t) \log(1 + |x|) \in L^1$ for all $0 < t < T$.

Proof. By (3.4) and (3.3), we have

$$\|B_1((m, v), (n, u))\|_{4/3} \lesssim \int_0^t (t-s)^{-\frac{1}{2} + \frac{3}{4} - 1} \|m(s) \nabla c(s) + m(s) u(s)\|_1 ds \\ \lesssim \int_0^t (t-s)^{-\frac{3}{4}} \|m(s)\|_{4/3} \|\nabla c(s) + u(s)\|_4 ds \\ \lesssim \left(\int_0^t (t-s)^{-\frac{3}{4}} s^{-\frac{1}{4}} s^{-\frac{1}{4}} ds \right) \|(m, v)\|_{X_T} \|(n, u)\|_{X_T} \\ \lesssim t^{-1/4} \|(m, v)\|_{X_T} \|(n, u)\|_{X_T}$$

and

$$\|B_2((m, v), (n, u))\|_4 \lesssim \int_0^t (t-s)^{-\frac{1}{2} + \frac{1}{4} - \frac{1}{2}} \|\nabla b(s) \otimes \nabla c(s) + v(s) \otimes u(s)\|_2 ds \\ \lesssim \int_0^t (t-s)^{-\frac{3}{4}} (\|\nabla b(s)\|_4 \|\nabla c(s)\|_4 + \|v(s)\|_4 \|u(s)\|_4) ds \\ \lesssim \int_0^t (t-s)^{-\frac{3}{4}} \left(\|m(s)\|_{4/3} \|n(s)\|_{4/3} + \|v(s)\|_4 \|u(s)\|_4 \right) ds \\ \lesssim \left(\int_0^t (t-s)^{-\frac{3}{4}} s^{-\frac{1}{4}} s^{-\frac{1}{4}} ds \right) \|(m, v)\|_{X_T} \|(n, u)\|_{X_T} \\ \lesssim t^{-1/4} \|(m, v)\|_{X_T} \|(n, u)\|_{X_T}.$$

Thus, we have proved that

$$\|B((m, v), (n, u))\|_{X_T} \lesssim \|(m, v)\|_{X_T} \|(n, u)\|_{X_T}. \quad (3.5)$$

We claim that if $n_0 \in L^1$ and $u_0 \in L^2$, then

$$\lim_{t \rightarrow 0} t^{1/4} \left(\|e^{t\Delta} n_0\|_{4/3} + \|e^{t\Delta} u_0\|_4 \right) = 0.$$

The limit holds if n_0 and u_0 are smooth and have a compact support. Therefore, the claim holds by a density argument. With this claim and the bilinear estimate (3.5), the local existence of a solution $(n, u) \in X_T$ of (1.1) follows from the Picard iteration. The solution (n, u) is in Y_T , and hence a mild solution of (1.1), since

$$\|n(t)\|_1 = \int_{\mathbb{R}^2} n(x, t) dx = \int_{\mathbb{R}^2} e^{t\Delta} n_0(x) dx = \int_{\mathbb{R}^2} n_0(x) dx$$

and

$$\|u(t)\|_2 \leq \|e^{t\Delta} u_0\|_2 + \|B_2((n, u), (n, u))\|_2 \lesssim \|u_0\|_2 + \|(n, u)\|_{X_T}^2.$$

It is easy to see

$$\|n(t) - n_0\|_1 \leq \|e^{t\Delta} n_0 - n_0\|_1 + \left\| \int_0^t e^{(t-s)\Delta} \nabla \cdot (n(s) \nabla c(s) + n(s) u(s)) ds \right\|_1 \rightarrow 0$$

and

$$\|u(t) - u_0\|_2 \leq \|e^{t\Delta} u_0 - u_0\|_2 + \left\| \int_0^t e^{(t-s)\Delta} \mathbf{P} \nabla \cdot (\nabla c(s) \otimes \nabla c(s) + u(s) \otimes u(s)) ds \right\|_1 \rightarrow 0,$$

as $t \rightarrow 0_+$, which proves (i).

From (3.1) and (3.2), using (3.3), we have for $1 \leq q \leq p \leq \infty$ ($p \neq 1, q \neq \infty$) that

$$\begin{aligned} \|n(t)\|_p &\lesssim \|e^{t\Delta} n_0\|_p + \int_0^t (t-s)^{-\frac{1}{2} + \frac{1}{p} - \frac{1}{q}} \|n(s) \nabla c(s) + n(s) u(s)\|_q ds \\ &\lesssim t^{\frac{1}{p}-1} \|n_0\|_1 + \int_0^t (t-s)^{\frac{1}{p} - \frac{2}{a}} \left(\|n(s)\|_a^2 + \|n(s)\|_a \|u(s)\|_{\frac{2a}{2-a}} \right) ds \end{aligned} \quad (3.6)$$

and for $1 \leq r \leq \frac{2p}{2-p} \leq \infty$ ($\frac{2p}{2-p} \neq 1, r \neq \infty$) that

$$\begin{aligned} \|u(t)\|_{\frac{2p}{2-p}} &\lesssim \|e^{t\Delta} u_0\|_{\frac{2p}{2-p}} + \int_0^t (t-s)^{-\frac{1}{2} + \frac{2-p}{2p} - \frac{1}{r}} \|\nabla c(s) \otimes \nabla c(s) + u(s) \otimes u(s)\|_r ds \\ &\lesssim t^{\frac{1}{p}-1} \|u_0\|_2 + \int_0^t (t-s)^{\frac{1}{p} - \frac{2}{a}} \left(\|n(s)\|_a^2 + \|u(s)\|_{\frac{2a}{2-a}}^2 \right) ds \end{aligned} \quad (3.7)$$

where $a = \frac{4q}{2+q} = \frac{2r}{1+r} < 2$ and $\frac{1}{p} - \frac{2}{a} > -1$.

For $1 \leq p < 2$, we have $1 \leq \frac{2p}{2-p} < \infty$. Taking $a = \frac{4}{3}$ so that $q = 1$, $r = 2$, and $\frac{1}{p} - \frac{2}{a} > -1$, (3.6) and (3.7) becomes

$$\|n(t)\|_p + \|u(t)\|_{\frac{2p}{2-p}} \lesssim t^{\frac{1}{p}-1} + \int_0^t (t-s)^{\frac{1}{p} - \frac{3}{2}} s^{-\frac{1}{2}} ds \lesssim t^{\frac{1}{p}-1} \quad \forall 1 \leq p < 2, \quad (3.8)$$

which completes the proof of (iii).

For $p = 2$, we choose $a \in (\frac{4}{3}, 2)$ in (3.6) and (3.7), say $a = \frac{5}{3}$ so that $q = \frac{10}{7}$, $r = 5$, and $\frac{1}{p} - \frac{2}{a} = -\frac{7}{10} > -1$, and get

$$\|n(t)\|_2 + \|u(t)\|_\infty \lesssim t^{-\frac{1}{2}} + \int_0^t (t-s)^{-\frac{7}{10}} s^{-\frac{4}{5}} ds \lesssim t^{-\frac{1}{2}},$$

where we used (3.8) with $p = \frac{5}{3}$.

For $2 < p < \infty$, we choose $a = \frac{4p}{1+2p} \in (\frac{8}{5}, 2)$ in (3.6) so that $q = \frac{2p}{1+p} \leq p$, and $\frac{1}{p} - \frac{2}{a} = \frac{1-2p}{2p} > -1$, and get

$$\|n(t)\|_p \lesssim t^{\frac{1}{p}-1} + \int_0^t (t-s)^{-1+\frac{1}{2p}} s^{\frac{1}{2p}-1} ds \lesssim t^{\frac{1}{p}-1} \quad \forall 2 < p < \infty,$$

where we used (3.8) with $p \in (\frac{8}{5}, 2)$.

For $p = \infty$, from (2.8), $\|\nabla c\|_\infty \lesssim \|n\|_1^{\frac{1}{4}} \|n\|_3^{\frac{3}{4}}$. Therefore,

$$\begin{aligned} \|n(t)\|_\infty &\lesssim t^{-1} \|n(t/2)\|_1 + \int_{t/2}^t (t-s)^{-\frac{1}{2}-\frac{1}{3}} (\|\nabla c(s)\|_\infty + \|u(s)\|_\infty) \|n(s)\|_3 ds \\ &\lesssim t^{-1} \|n_0\|_1 + \int_{t/2}^t (t-s)^{-\frac{5}{6}} s^{-\frac{7}{6}} ds \\ &\lesssim t^{-1}, \end{aligned}$$

proving (ii).

We omit the proof of (iv) and (v) as they follow from the technique used in the proof of [66, Proposition 2.3]. The positivity of mild solutions (vi) is a direct consequence of the strong maximum principle. In fact, the proof is identical to that of [66, Proposition 2.7] for PKS system in which the n -equation is tested against n^- , the negative part of the density n . In our case, when we test (1.1)₁ against n^- , the additional term vanishes because $\int_{\mathbb{R}^2} u \cdot \nabla n \cdot n^- dx = \int_{\mathbb{R}^2} u \cdot \nabla n^- \cdot n^- dx = 0$. We skip the full details and refer the interested reader to the proof of [66, Proposition 2.7].

To show (vii), note that

$$|\nabla \log(1 + |x|^2)| \leq \frac{2|x|}{1 + |x|^2} \leq 1 \quad \text{and} \quad |\Delta \log(1 + |x|^2)| = \frac{4}{(1 + |x|^2)^2} \leq 4.$$

So

$$\begin{aligned} &\frac{d}{dt} \int_{\mathbb{R}^2} n \log(1 + |x|^2) dx \\ &= - \int_{\mathbb{R}^2} u \cdot \nabla n \log(1 + |x|^2) dx + \int_{\mathbb{R}^2} \Delta n \log(1 + |x|^2) dx - \int_{\mathbb{R}^2} \nabla \cdot (n \nabla c) \log(1 + |x|^2) dx \\ &= \int_{\mathbb{R}^2} n u \cdot \nabla \log(1 + |x|^2) dx + \int_{\mathbb{R}^2} n \Delta \log(1 + |x|^2) dx + \int_{\mathbb{R}^2} n \nabla c \cdot \nabla \log(1 + |x|^2) dx \\ &\leq \int_{\mathbb{R}^2} n |u| dx + 4M + \int_{\mathbb{R}^2} n |\nabla c| dx \\ &\leq \|n\|_{4/3} \|u\|_4 + 4M + \|n\|_{4/3} \|\nabla c\|_4 \\ &\leq 4M + C(\|n_0\|_1, \|u_0\|_2) t^{-\frac{1}{2}}. \end{aligned}$$

Therefore

$$\int_{\mathbb{R}^2} n(t) \log(1 + |x|^2) dx \leq \int_{\mathbb{R}^2} n_0 \log(1 + |x|^2) dx + 4Mt + C(\|n_0\|_1, \|u_0\|_2) t^{\frac{1}{2}}. \quad (3.9)$$

□

Remark 3.1. It follows the same argument as in [66, Remark 2.1] with the estimate of bilinear forms B_1 and B_2 that if $n_0 \in L^1 \cap L^p$ and $u_0 \in L^2 \cap L^{\frac{2p}{2-p}}$ for $4/3 \leq p < 2$, then $n \in BC([0, T]; L^1 \cap L^p)$ and $u \in BC([0, T]; L^2 \cap L^{\frac{2p}{2-p}})$.

Remark 3.2. By Lemma 1.1, the mild solution constructed in Theorem 3.2 is a free-energy solution because it is a classical solution.

In order to have a better control of the modified free energy $H[n, u]$ in Section 4, we derive the following energy estimate of u using the free energy $\mathcal{F}[n, u]$ and the logarithmic Hardy–Littlewood–Sobolev inequality.

Lemma 3.3 (A priori bound of u). *Suppose that (n_0, u_0) with n_0 satisfying (1.3) and $u_0 \in L^2_\sigma$, and $\|n_0\|_1 \leq 8\pi$. Let (n, u) be a mild solution on $[0, T)$ given in Theorem 3.2. Then*

$$\frac{1}{2} \|u(t)\|_2^2 + \int_0^t \|\nabla u(\tau)\|_2^2 d\tau \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T).$$

The constant C is independent of T if $\|n_0\|_1 = 8\pi$.

Proof. By Lemma 1.1,

$$\mathcal{F}[n, u](t) = \mathcal{F}[n_0, u_0] - \int_0^t \int_{\mathbb{R}^2} [n|\nabla(\log n - c)|^2 + |\nabla u|^2] dx.$$

By Proposition 2.4,

$$\mathcal{F}[n_0, u_0] \geq \left(1 - \frac{M}{8\pi}\right) \int_{\mathbb{R}^2} n(t) \log n(t) dx - \frac{M}{8\pi} C(M) + \frac{1}{2} \|u(t)\|_2^2 + \int_0^t \|\nabla u(\tau)\|_2^2 d\tau,$$

or

$$\frac{1}{2} \|u(t)\|_2^2 + \int_0^t \|\nabla u(\tau)\|_2^2 d\tau \leq \mathcal{F}[n_0, u_0] - \left(1 - \frac{M}{8\pi}\right) \int_{\mathbb{R}^2} n(t) \log n(t) dx + \frac{M}{8\pi} C(M).$$

Our goal is to bound $\int_{\mathbb{R}^2} n \log n$ from below. In view of Proposition 2.4, it suffices to bound $\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} n(x)n(y) \log|x-y| dx dy$ from above. In fact, since

$$\begin{aligned} |c(x)| &\leq \frac{1}{2\pi} \int_{|x-y|\leq 1} n(y) \log \frac{1}{|x-y|} dy + \frac{1}{2\pi} \int_{|x-y|\geq 1} n(y) \log|x-y| dy \\ &\leq \frac{1}{2\pi} \int_{|x-y|\leq 1} \left[n(y) \log(1+n(y)) + \frac{C}{|x-y|} \right] dy \\ &\quad + \frac{1}{2\pi} \int_{|x-y|\geq 1} n(y) [\log(1+|x|) + \log(1+|y|)] dy \\ &\lesssim 1 + \log(1+|x|), \end{aligned}$$

we have

$$\int_{\mathbb{R}^2} \int_{\mathbb{R}^2} n(x)n(y) \log|x-y| dx dy = -2\pi \int_{\mathbb{R}^2} n(x)c(x) dx \lesssim 1 + \int_{\mathbb{R}^2} n(x) \log(1+|x|^2) dx.$$

Thus, we have

$$\begin{aligned} \int_{\mathbb{R}^2} n(t) \log n(t) dx &\geq -C(M) - \frac{2}{M} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} n(x, t)n(y, t) \log|x-y| dx dy \\ &\geq -C(M) - \int_{\mathbb{R}^2} n(x) \log(1+|x|^2) dx \\ &\geq -C(M) - \int_{\mathbb{R}^2} n_0 \log(1+|x|^2) dx - 4Mt - C(\|n_0\|_1, \|u_0\|_2) t^{\frac{1}{2}}, \end{aligned}$$

where we used (3.9) in the last inequality. We conclude that

$$\begin{aligned}
& \frac{1}{2} \|u(t)\|_2^2 + \int_0^t \|\nabla u(\tau)\|_2^2 d\tau \\
& \leq \mathcal{F}[n_0, u_0] - \left(1 - \frac{M}{8\pi}\right) \int_{\mathbb{R}^2} n(t) \log n(t) dx + \frac{M}{8\pi} C(M) \\
& \leq \mathcal{F}[n_0, u_0] + \left(1 - \frac{M}{8\pi}\right) \left[C(M) + \|n_0 \log(1 + |x|^2)\|_1 + 4MT + C(\|n_0\|_1, \|u_0\|_2) T^{\frac{1}{2}} \right] \\
& \quad + \frac{M}{8\pi} C(M),
\end{aligned}$$

completing the proof. \square

Remark 3.3. By [76, Lemma V.1.2.1 b)]

$$\left(\int_0^T \|u(\tau)\|_q^s d\tau \right)^{1/s} \lesssim \frac{1}{2} \left(\operatorname{ess\,sup}_{t \in [0, T]} \|u(t)\|_2 \right)^2 + \int_0^T \|\nabla u(\tau)\|_2^2 d\tau$$

for $2 \leq q < \infty$, $1 \leq s \leq \infty$ satisfying $\frac{2}{q} + \frac{2}{s} = 1$. Therefore, Lemma 3.3 gives

$$u \in L^s(0, T; L^q(\mathbb{R}^2)), \quad 2 \leq q < \infty, \quad 1 \leq s \leq \infty, \quad \frac{2}{q} + \frac{2}{s} = 1$$

with

$$\|u\|_{L^s(0, T; L^q(\mathbb{R}^2))} \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T)$$

in which the constant C is independent of T if $\|n_0\|_1 = 8\pi$.

Remark 3.4. If, additionally, $u_0 \in W_{0, \sigma}^{1,2}(\mathbb{R}^2)$ then $\partial_t u$, $u \cdot \nabla u$ and ∇P are in $L^2(\mathbb{R}^2 \times (0, T))$ by [76, Theorem V.1.8.1]. Note that [76, Theorem V.1.8.1] holds for the case $f = \nabla \cdot F$, $F \in L^4(\mathbb{R}^2 \times (0, T))$.

4. A PRIORI ENTROPY ESTIMATE FOR THE CAUCHY PROBLEM OF THE ADVECTION-PATLAK-KELLER-SEGEL SYSTEM (1.1)₁-(1.1)₂

This section is devoted to prove Proposition 1.4, following a similar approach as in [68] for PKS system. The key to derive the global-in-time a priori entropy estimate of (1.1)₁-(1.1)₂ is to split \mathbb{R}^2 into a ball (interior region), the complement of a larger ball (exterior region), and the annulus that connects the two regions. The main difficulties here for the PKS-NS system come from the coupling of velocity u which is not present in [68], and extra efforts are needed to control the terms generated from u . We will use the regularity estimates of Navier-Stokes equation with forcing.

4.1. Entropy estimates for exterior regions. In this subsection, we obtain the a priori estimate for large $|x|$. To this end, we set

$$H_{\text{ext}}(t; R) := \int_{|x| \geq R} [(1+n(t)) \log(1+n(t)) - n(t)] dx,$$

and introduce the Littlewood–Paley partition of unity.

Definition 4.1. Let $\phi(r)$ be a function in $C_0^\infty(0, \infty)$ such that $\phi(r) \in [0, 1]$ and $\text{supp } \phi \subset (1/2, 2)$. Then a family of the cut off functions $\{\Phi_k\}_{k=0}^\infty$ is called as the Littlewood–Paley partition of the unity in \mathbb{R}^2 if they satisfy

$$\Phi_k(x) = \phi(2^{-k}|x|), \quad x \in \mathbb{R}^2, k = 0, 1, 2, \dots$$

and $\sum_{k=0}^\infty \Phi_k \in C^\infty(0, \infty)$ with $\sum_{k=0}^\infty \Phi_k(x) = 1$ for $|x| > 1$ and $\text{supp } (\sum_{k=0}^\infty \Phi_k) \subset (1/2, \infty)$.

The Littlewood–Paley partition of unity $\{\Phi_k\}_{k=0}^\infty$ satisfies $\Phi_k \in C_0^\infty(\mathbb{R}^2)$, $\Phi_k(x) \in [0, 1]$,

$$\text{supp } \Phi_k \subset A_k := \{x \in \mathbb{R}^2 : 2^{k-1} \leq |x| \leq 2^{k+1}\},$$

and

$$|\nabla \Phi_k| \leq C 2^{-k} \Phi_k^{5/6}, \quad |\nabla \Phi_k^{1/2}| \leq C 2^{-k} \Phi_k^{1/3}, \quad |\nabla^2 \Phi_k| \leq C 2^{-2k} \Phi_k^{2/3},$$

where $C > 0$ is a constant independent of k .

Proposition 4.2. There exists a sufficiently large R_0 depending only on T , $\|n_0\|_1$, and $\|u_0\|_2$ such that

$$\begin{aligned} \sup_{0 \leq t < T} H_{\text{ext}}(t; R_0) + \int_0^T \int_{|x| \geq R_0} \frac{|\nabla n(t)|^2}{1+n(t)} dx dt \\ \leq 2H_{\text{ext}}(t; R_0/2) + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2)T, \end{aligned} \quad (4.1)$$

$$\int_0^T \int_{|x| \geq R_0} n^2(t) dx dt \leq C \|n_0\|_1 H_{\text{ext}}(t; R_0/2) + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2)T. \quad (4.2)$$

Proof. For fixed k , we let $c_m(t) = c(t) - [c(t)]_B$, where $[c(t)]_B = (1/|B|) \int_B c(t) dx$, $B = B_{2^{k+2}}(0)$. The equation exchanging $c(t)$ into $c_m(t)$ also holds, that is, $-\Delta c_m = n$. Note that by the equation of n ,

$$\partial_t \{[(1+n) \log(1+n) - n] \Phi_k\} = -u \cdot \nabla n \log(1+n) \Phi_k + (\Delta n) \log(1+n) \Phi_k - \nabla \cdot (n \nabla c) \log(1+n) \Phi_k.$$

Since

$$\begin{aligned} (\Delta n) \log(1+n) \Phi_k &= \nabla \cdot [\nabla n \log(1+n) \Phi_k - ((1+n) \log(1+n) - n) \nabla \Phi_k] \\ &\quad + ((1+n) \log(1+n) - n) \Delta \Phi_k - \frac{|\nabla n|^2}{1+n} \Phi_k \end{aligned}$$

and

$$\begin{aligned} -\nabla \cdot (n \nabla c_m) \log(1+n) \Phi_k &= -\nabla \cdot [n \log(1+n) (\nabla c_m) \Phi_k] + n \log(1+n) (\nabla c_m) \cdot \nabla \Phi_k \\ &\quad + n \nabla \log(1+n) \cdot (\nabla c_m) \Phi_k, \end{aligned}$$

we have

$$\begin{aligned} \partial_t \{[(1+n) \log(1+n) - n] \Phi_k\} &+ \frac{|\nabla n|^2}{1+n} \Phi_k \\ &= -u \cdot \nabla n \log(1+n) \Phi_k \\ &\quad + \nabla \cdot \{ \nabla n \log(1+n) \Phi_k - [(1+n) \log(1+n) - n] \nabla \Phi_k - n \log(1+n) (\nabla c_m) \Phi_k \} \\ &\quad + [(1+n) \log(1+n) - n] \Delta \Phi_k + n \log(1+n) \nabla c_m \cdot \nabla \Phi_k \\ &\quad + n \nabla \log(1+n) \cdot (\nabla c_m) \Phi_k. \end{aligned}$$

We then have

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^2} \{[(1+n) \log(1+n) - n] \Phi_k\} dx + \int_{\mathbb{R}^2} \frac{|\nabla n|^2}{1+n} \Phi_k dx \\ = - \int_{\mathbb{R}^2} u \cdot \nabla n \log(1+n) \Phi_k dx + I + II + III + IV, \end{aligned}$$

where

$$\begin{aligned} |I| + |II| + |III| + |IV| \\ \leq \left(9\varepsilon + C \int_{A_k} n(t) dx\right) \int_{\mathbb{R}^2} \frac{|\nabla n(t)|^2}{1+n(t)} \Phi_k dx \\ + C2^{-2n} (\|n_0\|_1 + \|n_0\|_1^2) + C(1+\varepsilon)^2 \int_{\mathbb{R}^2} n(t) \Phi_k dx \\ + C\varepsilon^{-5} 2^{-4n} \|c(t)\|_{\text{BMO}}^6 + C2^{-4n} \|c(t)\|_{\text{BMO}}^3 + C2^{-4n}, \end{aligned}$$

where ε is an arbitrary number with $0 < \varepsilon < 1$ (see [68, (3.20)]). Now, using the fact that $\nabla \cdot u = 0$ and the similar computation for IV on page 87 of [68], we get

$$\begin{aligned} - \int_{\mathbb{R}^2} u \cdot \nabla n \log(1+n) \Phi_k dx \\ = - \int_{\mathbb{R}^2} u \cdot \nabla [(1+n) \log(1+n) - n] \Phi_k dx \\ = \int_{\mathbb{R}^2} u \cdot \nabla \Phi_k [(1+n) \log(1+n) - n] dx \\ \leq C2^{-k} \int_{A_k} |u| [(1+n) \log(1+n) - n] \Phi_k^{5/6} dx \\ \leq C2^{-k} \left(\int_{A_k} [(1+n) \log(1+n) - n]^{3/2} \Phi_k^{5/4} dx \right)^{2/3} \left(\int_{A_k} |u|^3 dx \right)^{1/3} \\ \leq \int_{A_k} [(1+n) \log(1+n) - n]^{3/2} \Phi_k dx + C2^{-3k} \|u\|_3^3 \\ \leq \left(\int_{A_k} n(t) dx \right) \left(\int_{\mathbb{R}^2} \frac{|\nabla n(t)|^2}{1+n(t)} \Phi_k dx \right) \\ + C\varepsilon 2^{-2k} \left(\int_{\mathbb{R}^2} n_0 dx \right)^2 + C\varepsilon(1+\varepsilon) \int_{\mathbb{R}^2} n(t) \Phi_k dx + C2^{-3k} \|u\|_3^3. \end{aligned}$$

By the same argument as in the proof of [68, Proposition 3.2] and taking $\varepsilon = 1/36$, we can choose N sufficiently large such that for all $k \geq N$,

$$9\varepsilon + (C+1) \sup_{0 \leq t < T} \int_{A_k} n(t) dx \leq \frac{1}{2},$$

and thus

$$\begin{aligned} \frac{d}{dt} \int_{\mathbb{R}^2} \{[(1+n) \log(1+n) - n] \Phi_k\} dx + \frac{1}{2} \int_{\mathbb{R}^2} \frac{|\nabla n|^2}{1+n} \Phi_k dx \\ \leq C(1+\varepsilon)^2 \int_{\mathbb{R}^2} n(t) \Phi_k dx + C2^{-2k} \left(\|c(t)\|_{\text{BMO}}^6 + \|c(t)\|_{\text{BMO}}^3 + \|c(t)\|_{\text{BMO}}^2 + \|c(t)\|_{\text{BMO}} + 1 \right) \\ + C2^{-3k} \|u\|_3^3 \\ \leq C(1+\varepsilon)^2 \int_{\mathbb{R}^2} n(t) \Phi_k dx + C2^{-2k} \left(\|n_0\|_1^6 + \|n_0\|_1^3 + \|n_0\|_1^2 + \|n_0\|_1 + 1 \right) + C2^{-3k} \|u\|_3^3 \end{aligned}$$

since $\|c(t)\|_{\text{BMO}} \lesssim \|n(t)\|_1 = \|n_0\|_1$ by Proposition 2.2. Thus, we have

$$\begin{aligned} & \int_{\mathbb{R}^2} [(1+n(t)) \log(1+n(t)) - n(t)] \Phi_k dx + \frac{1}{2} \int_0^t \int_{\mathbb{R}^2} \frac{|\nabla n|^2}{1+n} \Phi_k dx ds \\ & \leq \int_{\mathbb{R}^2} [(1+n_0) \log(1+n_0) - n_0] \Phi_k dx + C \int_0^T \int_{\mathbb{R}^2} n(t) \Phi_k dx \\ & \quad + C2^{-2k} \left(\|n_0\|_1^6 + \|n_0\|_1^3 + \|n_0\|_1^2 + \|n_0\|_1 + 1 \right) T + C2^{-3k} \int_0^T \|u(t)\|_3^3 dt. \end{aligned} \quad (4.3)$$

Note that we have $\|u\|_{L^6(0,T;L^3(\mathbb{R}^2))} \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2)$ by Remark 3.3. Then

$$\int_0^T \|u(t)\|_3^3 dt \leq \int_0^T \left(\frac{\|u(t)\|_3^6}{2} + \frac{1}{2} \right) dt \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2) T. \quad (4.4)$$

Adding (4.3) for $k \geq N$ and using $\sum_{k=N}^{\infty} \Phi_k(x) = 1$ for $|x| \geq 2^N$, we have

$$\begin{aligned} & \int_{|x| \geq 2^N} [(1+n(t)) \log(1+n(t)) - n(t)] dx + \frac{1}{2} \int_0^t \int_{|x| \geq 2^N} \frac{|\nabla n|^2}{1+n} \Phi_k dx ds \\ & \leq \int_{|x| \geq 2^{N-1}} [(1+n_0) \log(1+n_0) - n_0] dx + C \left(\|n_0\|_1^6 + \|n_0\|_1^3 + \|n_0\|_1^2 + \|n_0\|_1 + 1 \right) T \\ & \quad + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2) T, \end{aligned}$$

proving (4.1) with $R_0 = 2^N$. Next, we have (see [68, (3.24)])

$$\int_{\mathbb{R}^2} n^2(t) \Phi_k dx \leq 2 \|n_0\|_1 \left(\int_{\mathbb{R}^2} \frac{|\nabla n|^2}{1+n} \Phi_k dx \right) + C2^{-2k} \|n_0\|_1^2 + 4 \int_{\mathbb{R}^2} n(t) \Phi_k dx. \quad (4.5)$$

Using (4.3) to the first term on the right hand side of (4.5) and adding these for all $k \geq N$, we conclude that

$$\begin{aligned} & \int_0^T \int_{|x| \geq 2^N} n^2 dx dt \leq 2 \|n_0\|_1 \int_0^T \int_{|x| \geq 2^N} \frac{|\nabla n|^2}{1+n} dx dt + C(\|n_0\|_1^2 + \|n_0\|_1) T \\ & \leq 4 \|n_0\|_1 \int_{|x| \geq 2^{N-1}} [(1+n_0) \log(1+n_0) - n_0] dx \\ & \quad + C \|n_0\|_1 \left(\|n_0\|_1^6 + \|n_0\|_1^3 + \|n_0\|_1^2 + \|n_0\|_1 + 1 \right) T \\ & \quad + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2) \|n_0\|_1 T + C(\|n_0\|_1^2 + \|n_0\|_1) T. \end{aligned}$$

This proves (4.2). \square

4.2. A priori estimates for exterior regions. Let $R_0 > 0$ be given in Proposition 4.2, and choose it larger so that $n_0(x) \leq 1$ for $|x| > R_0$. Then

$$\begin{aligned} & \sup_{0 < t < T} \int_{|x| \geq R_0} (1+n(t)) \log(1+n(t)) dx \leq C(n_0, u_0, T), \\ & \int_0^T \int_{|x| \geq R_0} \frac{|\nabla n|^2}{1+n} dx dt \leq C(n_0, u_0, T), \\ & \int_0^T \int_{|x| \geq R_0} n^2 dx dt \leq C(n_0, u_0, T), \end{aligned}$$

and for $R \geq R_0$ and $p \in [1, \infty)$

$$\int_{|x| \geq R} n_0^p(x) dx \leq \int_{|x| \geq R} n_0(x) dx.$$

For $R > 1$, let $\Phi_R \in C^\infty(\mathbb{R}^2)$ be such that $\Phi_R(x) \in [0, 1]$, $\Phi_R(x) = 1$ for $|x| \geq R$, $\text{supp } \Phi_R \subset \mathbb{R}^2 \setminus \overline{B_{R/2}(0)}$, and

$$|\nabla \Phi_R| \leq CR^{-1} \Phi_R^{5/6}, \quad |\nabla(\Phi_R^{1/2})| \leq CR^{-1} \Phi_R^{1/3}, \quad |\nabla^2 \Phi_R| \leq CR^{-2} \Phi_R^{2/3},$$

where $C > 0$ is a constant independent of R . Then

$$\text{supp}(\nabla \Phi_R) \subset A_R^* := \{x \in \mathbb{R}^2 : R/2 \leq |x| \leq R\}.$$

Lemma 4.3. *For any $R \geq 2R_0$,*

$$\sup_{0 < t < T} \int_{|x| \geq R} n^2(t) dx + \frac{1}{2} \int_0^T \int_{|x| \geq R} |\nabla n|^2 dx dt \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R). \quad (4.6)$$

Proof. For fixed $R \geq 2R_0$, we let $c_m(t) = c(t) - [c(t)]_B$, where $[c(t)]_B = (1/|B|) \int_B c(t) dx$, $B = B_{2R}(0)$. Then $\|c_m\|_p \lesssim \|c\|_{BMO}$ for $1 < p < \infty$ by Lemma 2.3. Multiplying the equation of n by $n\Phi_R$, integrating on \mathbb{R}^2 , and performing integration by parts, we have

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^2} n^2(t) \Phi_R dx + \int_{\mathbb{R}^2} |\nabla n(t)|^2 \Phi_R dx \\ &= - \int_{\mathbb{R}^2} nu \cdot \nabla n \Phi_R dx + \int_{\mathbb{R}^2} n(\Delta n) \Phi_R dx - \int_{\mathbb{R}^2} n \nabla \cdot (n \nabla c) \Phi_R dx + \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx \\ &= - \int_{\mathbb{R}^2} nu \cdot \nabla n \Phi_R dx + \frac{1}{2} \int_{\mathbb{R}^2} n^3 \Phi_R dx + \frac{1}{2} \int_{\mathbb{R}^2} n^2(1 - c_m) \Delta \Phi_R dx - \int_{\mathbb{R}^2} n \nabla n \cdot \nabla \Phi_R c_m dx, \end{aligned}$$

where

$$\frac{1}{2} \int_{\mathbb{R}^2} n^2(1 - c_m) \Delta \Phi_R dx \leq \frac{1}{4} \int_{\mathbb{R}^2} n^3 \Phi_R dx + CR^{-4} (\|c\|_{BMO}^3 + 1)$$

and

$$- \int_{\mathbb{R}^2} n \nabla n \cdot \nabla \Phi_R c_m dx \leq \frac{1}{3} \int_{\mathbb{R}^2} n^3 \Phi_R dx + \frac{1}{2} \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx + CR^{-4} \|c\|_{BMO}^6.$$

Hence

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^2} n^2 \Phi_R dx + \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx \\ & \leq - \int_{\mathbb{R}^2} nu \cdot \nabla n \Phi_R dx + \frac{13}{6} \int_{\mathbb{R}^2} n^3 \Phi_R dx + CR^{-4} (\|c\|_{BMO}^6 + \|c\|_{BMO}^3 + 1), \end{aligned}$$

where

$$\begin{aligned} - \int_{\mathbb{R}^2} nu \cdot \nabla n \Phi_R dx &= - \frac{1}{2} \int_{\mathbb{R}^2} u \cdot \nabla(n^2) \Phi_R dx = \frac{1}{2} \int_{\mathbb{R}^2} n^2 u \cdot \nabla \Phi_R dx \\ &\leq \frac{1}{2} CR^{-1} \int_{A_R^*} n^2 |u| \Phi_R dx \\ &\leq CR^{-3} \|u\|_3^3 + \frac{5}{6} \int_{\mathbb{R}^2} n^3 \Phi_R dx \end{aligned}$$

since u is divergence-free. Thus,

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^2} n^2 \Phi_R dx + \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx \\ & \leq 3 \int_{\mathbb{R}^2} n^3 \Phi_R dx + CR^{-4} \left(\|c\|_{\text{BMO}}^6 + \|c\|_{\text{BMO}}^3 + 1 \right) + CR^{-3} \|u\|_3^3, \end{aligned}$$

and from a similar argument as that of [68, (3.31)] we have

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^2} n^2 \Phi_R dx + \left(1 - \varepsilon \int_{A_R^*} (1+n) \log(1+n) dx \right) \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx \\ & \leq C \left(\int_{\mathbb{R}^2} n^{3/2} |\nabla \Phi_R^{1/2}| dx \right)^2 + C(\varepsilon) \int_{\mathbb{R}^2} n \Phi_R dx + CR^{-4} \left(\|c\|_{\text{BMO}}^6 + \|c\|_{\text{BMO}}^3 + 1 \right) \\ & \quad + CR^{-3} \|u\|_3^3. \end{aligned}$$

Choosing $\varepsilon > 0$ such that

$$1 - \varepsilon \int_{A_R^*} (1+n) \log(1+n) dx = \frac{1}{2},$$

we have

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^2} n^2 \Phi_R dx + \frac{1}{2} \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx \\ & \leq CR^{-2} \|n_0\|_1 \int_{|x| \geq R/2} n^2 dx + C(\varepsilon) \|n_0\|_1 + C(\|n_0\|_1) R^{-4} + CR^{-3} \|u\|_3^3. \end{aligned}$$

Recall from (4.4) that $\int_0^T \|u(t)\|_3^3 dt \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2) T$. Integrating the above inequality from 0 to t with respect to the time variable, we have

$$\begin{aligned} \int_{\mathbb{R}^2} n^2(t) \Phi_R dx + \frac{1}{2} \int_0^t \int_{\mathbb{R}^2} |\nabla n|^2 \Phi_R dx ds & \leq \int_{\mathbb{R}^2} n_0^2 \Phi_R dx + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R) \\ & \leq \int_{\mathbb{R}^2} n_0 \Phi_R dx + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R). \end{aligned}$$

This proves (4.6). \square

Lemma 4.4. For any $R \geq 2^2 R_0$,

$$\int_0^T \int_{|x| \geq R} n^4 dx dt \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R).$$

Proof. The lemma is a consequence of Lemma 4.3 following the same proof of [68, Lemma 3.5]. \square

Lemma 4.5. For any $R \geq 2^3 R_0$,

$$\sup_{0 < t < T} \int_{|x| \geq R} n^3(t) dx + \int_0^T \int_{|x| \geq R} |\nabla n^{3/2}|^2 \Phi_R dx dt \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R).$$

Proof. Multiplying the equation of n by $n^2\Phi_R$, integrating over \mathbb{R}^2 , and using integration by parts and the fact that u is divergence-free, we have

$$\begin{aligned} & \frac{1}{3} \frac{d}{dt} \int_{\mathbb{R}^2} n^3 \Phi_R dx + \frac{8}{9} \int_{\mathbb{R}^2} |\nabla n^{3/2}|^2 \Phi_R dx \\ &= - \int_{\mathbb{R}^2} (u \cdot \nabla n) n^2 \Phi_R dx + \frac{1}{3} \int_{\mathbb{R}^2} n^3 \Delta \Phi_R dx - \int_{\mathbb{R}^2} n^2 \nabla \cdot (n \nabla c_m) \Phi_R dx \\ &= \frac{1}{3} \int_{\mathbb{R}^2} n^3 u \cdot \nabla \Phi_R dx + \frac{1}{3} \int_{\mathbb{R}^2} n^3 \Delta \Phi_R dx + \frac{1}{3} \int_{\mathbb{R}^2} n^3 \nabla c_m \cdot \nabla \Phi_R dx + \frac{2}{3} \int_{\mathbb{R}^2} n^4 \Phi_R dx, \end{aligned}$$

where $c_m(t) = c(t) - [c(t)]_B$, $[c(t)]_B = (1/|B|) \int_B c(t) dx$, $B = B_{2R}(0)$ satisfies $\|c_m\|_p \lesssim \|c\|_{BMO}$ by Lemma 2.3. So

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^2} n^3 \Phi_R dx + \frac{8}{9} \int_{\mathbb{R}^2} |\nabla n^{3/2}|^2 \Phi_R dx \\ &= \int_{A_R^*} n^3 u \cdot \nabla \Phi_R dx + \int_{\mathbb{R}^2} n^3 \Delta \Phi_R dx + \int_{\mathbb{R}^2} n^3 \nabla c_m \cdot \nabla \Phi_R dx + 2 \int_{\mathbb{R}^2} n^4 \Phi_R dx \\ &\leq CR^{-1} \int_{A_R^*} n^3 |u| \Phi_R^{5/6} dx + CR^{-2} \int_{|x| \geq R/2} n^3 dx + \int_{\mathbb{R}^2} n^3 \nabla c_m \cdot \nabla \Phi_R dx + 2 \int_{\mathbb{R}^2} n^4 dx \\ &\leq CR^{-1} \|u(t)\|_4^4 + CR^{-1} \int_{|x| \geq R/2} n^4 dx + F(t), \end{aligned}$$

where

$$\begin{aligned} F(t) &= CR^{-2} \int_{|x| \geq R/2} n^3 dx + C \int_{|x| \geq R/2} n^4 dx \\ &\quad + CR^{-6} \left(\|c(t)\|_{BMO}^8 + \|c(t)\|_{BMO}^4 \right), \end{aligned}$$

and $\|u\|_{L^4(\mathbb{R}^2 \times (0, T))} \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2)$ by Remark 3.3. By Lemma 4.3 and Lemma 4.4 we have

$$\int_0^T F(t) dx \leq C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R).$$

Therefore,

$$\begin{aligned} & \int_{\mathbb{R}^2} n^3(t) \Phi_R dx + \int_0^t \int_{\mathbb{R}^2} |\nabla n^{3/2}|^2 \Phi_R dx ds \\ &\leq \int_{\mathbb{R}^2} n_0^3 \Phi_R dx + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R) \\ &\leq \int_{\mathbb{R}^2} n_0 \Phi_R dx + C(\|n_0\|_1, \|n_0 \log n_0\|_1, \|u_0\|_2, T, R). \end{aligned}$$

This completes the proof of the lemma. \square

4.3. A priori estimates for annuli. Let $R_0 > 0$ be given in Proposition 4.2. Throughout the rest of this paper, we denote

$$A_R = \{x \in \mathbb{R}^2 : R/2 \leq |x| \leq 2R\}. \quad (4.7)$$

For $R \geq 1$ we let $\tilde{\Phi}_R \in C_0^\infty(\mathbb{R}^2)$ be such that $\tilde{\Phi}_R(x) \in [0, 1]$, $\tilde{\Phi}_R(x) = 1$ for $R/2 \leq |x| \leq 2R$,

$$\text{supp } \tilde{\Phi}_R \subset \tilde{A}_R := \{x \in \mathbb{R}^2 : R/3 \leq |x| \leq 3R\},$$

and

$$|\nabla\tilde{\Phi}_R| \leq CR^{-1}\tilde{\Phi}_R^{5/6}, \quad |\nabla\tilde{\Phi}_R^{1/2}| \leq CR^{-1}\tilde{\Phi}_R^{1/3}, \quad |\nabla^2\tilde{\Phi}_R| \leq CR^{-2}\tilde{\Phi}_R^{2/3},$$

where $C > 0$ is independent of R . We put $m(x, t) = n(x, t)\tilde{\Phi}_R(x)$, which satisfies $m \in C([0, T]; L^2(\mathbb{R}^2)) \cap L^\infty(0, T; L^2(\mathbb{R}^2))$ and

$$\partial_t m - \Delta m = f, \quad 0 < t < T, \quad x \in \mathbb{R}^2, \quad (4.8)$$

where

$$f = -u \cdot \nabla n \tilde{\Phi}_R - (2\nabla\tilde{\Phi}_R + \tilde{\Phi}_R \nabla c) \cdot \nabla n - (\Delta\tilde{\Phi}_R)n + \tilde{\Phi}_R n^2. \quad (4.9)$$

Denote $w(x, t) = u(x, t)\tilde{\Phi}_R(x)$, then

$$\partial_t w - \Delta w = g, \quad 0 < t < T, \quad x \in \mathbb{R}^2, \quad (4.10)$$

where

$$g = -(u \cdot \nabla)u\tilde{\Phi}_R - \nabla P\tilde{\Phi}_R + n\nabla c\tilde{\Phi}_R - 2\nabla u \cdot \nabla\tilde{\Phi}_R - u(\Delta\tilde{\Phi}_R). \quad (4.11)$$

Lemma 4.6. *There exists $R_1 \geq 2^5 R_0$ such that for any $R \geq R_1$, ∇c and c_m are bounded on $(0, T) \times A_R$, where $c_m(t) = c(t) - [c(t)]_B$ in which $[c(t)]_B = (1/|B|) \int_B c(t) dx$, $B = B_{2R}(0)$.*

Proof. The lemma is a consequence of Lemma 4.5 following the same proof of [68, Lemma 3.7]. \square

Lemma 4.7. *There exists $R_2 \geq 3R_1$ such that for any $R \geq R_2$, u is bounded on $A_R \times (0, T)$.*

Proof. By Remark 3.4 and Lemma 4.3, g defined by (4.11) is in $L^2(0, T; L^2(\mathbb{R}^2))$. The rest of the proof is identical to that of [68, Lemma 3.8]. \square

Lemma 4.8. *There exists $R_3 \geq 3R_2$ such that for any $R \geq R_3$, n is bounded on $A_R \times (0, T)$.*

Proof. By Lemmas in Subsection 4.2, Lemma 4.6 and Lemma 4.7, f defined by (4.9) is in $L^2(0, T; L^2(\mathbb{R}^2))$. The rest of the proof is identical to that of [68, Lemma 3.8]. \square

Lemma 4.9. *There exists $R_4 \geq 3^3 R_3$ such that for any $R \geq R_4$, ∇n is bounded on $A_R \times (0, T)$.*

Proof.

Step 1. We claim that

$$\sup_{0 < t < T} \|\nabla n(t)\|_{L^2(A_R)} < \infty. \quad (4.12)$$

Indeed, denoting $c_m(t) = c(t) - [c(t)]_B$, $B = B_{2R}(0)$, we have

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \int_{\mathbb{R}^2} |\nabla n|^2 \tilde{\Phi}_R dx \\
&= \int_{\mathbb{R}^2} u \cdot \nabla n (\Delta n) \tilde{\Phi}_R dx - \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx - \int_{\mathbb{R}^2} \nabla n \cdot \nabla c_m (\Delta n) \tilde{\Phi}_R dx - \int_{\mathbb{R}^2} n^2 \Delta n \tilde{\Phi}_R dx \\
&\quad + \int_{\mathbb{R}^2} (u \cdot \nabla n) (\nabla n \cdot \nabla \tilde{\Phi}_R) dx - \int_{\mathbb{R}^2} \Delta n \nabla n \cdot \nabla \tilde{\Phi}_R dx + \int_{\mathbb{R}^2} (\nabla n \cdot \nabla c_m) (\nabla n \cdot \nabla \tilde{\Phi}_R) dx \\
&\quad - \int_{\mathbb{R}^2} n^2 \nabla n \cdot \tilde{\Phi}_R dx \\
&\leq \int_{\mathbb{R}^2} u \cdot \nabla n (\Delta n) \tilde{\Phi}_R dx - \frac{3}{4} \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx + C \|\nabla c_m\|_{L^\infty(\tilde{A}_R)}^2 \int_{\tilde{A}_R} |\nabla n|^2 dx + C \int_{\tilde{A}_R} n^4 dx \\
&\quad + \int_{\mathbb{R}^2} (u \cdot \nabla n) (\nabla n \cdot \nabla \tilde{\Phi}_R) dx + \frac{1}{4} \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx + C (\|\nabla c_m\|_{L^\infty(\tilde{A}_R)} + 1) \int_{\tilde{A}_R} |\nabla n|^2 dx \\
&\quad + C \int_{\tilde{A}_R} n^4 dx.
\end{aligned}$$

Since

$$\begin{aligned}
& \int_{\mathbb{R}^2} u \cdot \nabla n (\Delta n) \tilde{\Phi}_R dx \\
&\leq \int_{\mathbb{R}^2} |u| |\nabla n| |\Delta n| \tilde{\Phi}_R dx \\
&\leq C(\varepsilon) \|u(t)\|_{L^\infty(\tilde{A}_R)} \int_{\tilde{A}_R} |\nabla n|^2 dx + \varepsilon \|u(t)\|_{L^\infty(\tilde{A}_R)} \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx \\
&\leq C(\varepsilon) \|u\|_{L^\infty(\tilde{A}_R \times (0, T))} \int_{\tilde{A}_R} |\nabla n|^2 dx + \varepsilon \|u\|_{L^\infty(\tilde{A}_R \times (0, T))} \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx \\
&\leq C(\varepsilon) \|u\|_{L^\infty(\tilde{A}_R \times (0, T))} \int_{\tilde{A}_R} |\nabla n|^2 dx + \frac{1}{4} \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx
\end{aligned}$$

by choosing $\varepsilon > 0$ sufficiently small, and

$$\begin{aligned}
\int_{\mathbb{R}^2} (u \cdot \nabla n) (\nabla n \cdot \nabla \tilde{\Phi}_R) dx &\leq CR^{-1} \int_{\mathbb{R}^2} |u| |\nabla n|^2 \tilde{\Phi}_R^{5/6} \\
&\leq CR^{-1} \|u(t)\|_{L^\infty(\tilde{A}_R)} \int_{\tilde{A}_R} |\nabla n|^2 \\
&\leq CR^{-1} \|u\|_{L^\infty(\tilde{A}_R \times (0, T))} \int_{\tilde{A}_R} |\nabla n|^2,
\end{aligned}$$

we obtain

$$\begin{aligned}
& \frac{d}{dt} \int_{\mathbb{R}^2} |\nabla n|^2 \tilde{\Phi}_R dx + \int_{\mathbb{R}^2} |\Delta n|^2 \tilde{\Phi}_R dx \\
&\leq C \left[\|\nabla c_m\|_{L^\infty(\tilde{A}_R)}^2 + \|\nabla c_m\|_{L^\infty(\tilde{A}_R)} + (C(\varepsilon) + CR^{-1}) \|u\|_{L^\infty(\tilde{A}_R \times (0, T))} + 1 \right] \int_{\tilde{A}_R} |\nabla n|^2 dx \\
&\quad + C \int_{\tilde{A}_R} n^4 dx.
\end{aligned}$$

The claim of **Step 1**, (4.12) is a direct consequence of the boundedness of ∇c , n and u in the annulus (Lemma 4.6, Lemma 4.7, Lemma 4.8) and a use of Grönwall's inequality.

Step 2. Let f be as in (4.9). By (4.12) and

$$\begin{aligned} \left\| (u \cdot \nabla n) \tilde{\Phi}_R \right\|_2 &\leq \|u(t)\|_{L^\infty(\tilde{A}_R)} \left(\int_{\mathbb{R}^2} |\nabla n|^2 \tilde{\Phi}_R dx \right)^{1/2} \\ &\leq \|u\|_{L^\infty(\tilde{A}_R \times (0, T))} \left(\int_{\mathbb{R}^2} |\nabla n|^2 \tilde{\Phi}_R dx \right)^{1/2}, \end{aligned}$$

we have

$$\sup_{0 < t < T} \|f(t)\|_2 < \infty$$

in view of Lemma 4.7. The rest of the proof is identical to that of [68, Lemma 3.9]. \square

Lemma 4.10. *There exists $R_5 \geq 3R_4$ such that for any $R \geq R_5$, $\nabla^2 c$ is bounded on $A_R \times (0, T)$.*

Proof. The lemma is a consequence of Lemma 4.8 and Lemma 4.9 following the same proof of [68, Lemma 3.10]. \square

Lemma 4.11. *There exists $R_6 \geq 3^2 R_5$ such that for any $R \geq R_6$, $\nabla^2 n$ is bounded on $A_R \times (0, T)$.*

Proof. Let f be as in (4.9). Since $\sup_{0 < t < T} \|u(t) \cdot \nabla n(t) \tilde{\Phi}_R\|_\infty < \infty$, we have

$$\sup_{0 < t < T} \|f(t)\|_\infty < \infty.$$

The rest of the proof is identical to that of [68, Lemma 3.11]. \square

Proposition 4.12. *There exists $\tilde{R}_0 > R_0$ such that for any $R \geq \tilde{R}_0$, the followings hold:*

- (i) $\partial_t^k \nabla_x^l n$ ($0 \leq 2k + l \leq 2$) are bounded on $A_R \times (0, T)$.
- (ii) $c_m, \nabla_x^l c$ ($1 \leq l \leq 2$), and u are bounded on $A_R \times (0, T)$.
- (iii) there exist x_0 satisfying $R < |x_0| < 2R$, and $\varepsilon_0 > 0, \delta > 0$ such that

$$n(x, t) \geq \delta \quad \text{for } |x - x_0| \leq \varepsilon_0, 0 \leq t < T.$$

Proof. The assertions (i) and (ii) follow from the preceding lemmas.

To prove (iii), we claim that $\nabla^l n$ ($0 \leq l \leq 2$) are uniformly Hölder continuous on $[T/2, T) \times \tilde{A}_R^*$, where $\tilde{A}_R^* := \{x \in \mathbb{R}^2 : 2R/3 \leq |x| \leq 8R/3\}$.

- (a) Since n satisfies $\partial_t n - \Delta n = -\nabla \cdot (n(u + \nabla c))$ and $n(u + \nabla c)$ is bounded on $(0, T) \times \{x \in \mathbb{R}^2 : R/3 < |x| < 3R\}$, applying local Schauder estimates for parabolic equations, we see that n is uniformly Hölder continuous on $[T/2, T) \times \tilde{A}_R^*$.
- (b) Similarly, by $\partial_t(\nabla n) - \Delta(\nabla n) = -\nabla \cdot (\nabla(n(u + \nabla c)))$ and the boundedness of $\nabla(n(u + \nabla c))$ in $(0, T) \times \{R/3 < |x| < 3R\}$, the uniform Hölder continuity of ∇n in $[T/2, T) \times \tilde{A}_R^*$ is deduced.
- (c) For $\nabla^2 n$, observe that

$$\partial_t(\partial_k \partial_j n) - \Delta(\partial_k \partial_j n) + \nabla c \cdot \nabla(\partial_k \partial_j n) - 2n(\partial_k \partial_j n) = \partial_k g_1 + g_2$$

where $g_1 = -\partial_j(u \cdot \nabla n) - \nabla n \cdot \nabla(\partial_j c)$ and $g_2 = -\nabla \partial_j n \cdot \nabla \partial_j c + 2\partial_k n \partial_j n$. Hence, by the boundedness of g_1 and g_2 on $(0, T) \times \{R/3 < |x| < 3R\}$, $\partial_k \partial_j n$ is uniformly Hölder continuous on $[T/2, T) \times \tilde{A}_R^*$.

The rest of the proof is identical to that of [68, Proposition 3.12] given that $u + \nabla c$ is bounded on $[T/2, T) \times \{x \in \mathbb{R}^2 : R < |x| < 2R\}$. \square

4.4. Entropy estimates for interior regions. We now derive the a priori estimate for small $|x|$. Define $\Psi_R(x) = \Psi(R^{-1}x)$ where $\Psi \in C^\infty(\mathbb{R}^2)$, $\Psi(x) \in [0, 1]$, $\Psi(x) = 1$ for $|x| \leq 1$, $\text{supp } \Psi \subset B_2(0)$. It follows from the definition of Ψ_R that

$$\text{supp } [\nabla \Psi_R] \subset A_R, \quad \text{supp } [\Delta \Psi_R] \subset A_R,$$

where A_R is an annulus defined in (4.7).

We define

$$\begin{aligned} H_{\text{int}}(t; R) &:= \int_{\mathbb{R}^2} [(1+n(t)) \log(1+n(t)) - n(t)] \Psi_R dx - \frac{1}{2} \int_{\mathbb{R}^2} n(t) c_m(t) \Psi_R dx \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^2} |u|^2 \Psi_R dx, \end{aligned}$$

where $c_m(t) = c(t) - [c(t)]_B$, $[c(t)]_B = (1/|B|) \int_B c(t) dx$, $B = B_{2R}(0)$.

Lemma 4.13.

$$\begin{aligned} \frac{d}{dt} H_{\text{int}}(t; R) + \int_{\mathbb{R}^2} n |\nabla \log(1+n(t)) - \nabla c_m(t)|^2 \Psi_R dx + \int_{\mathbb{R}^2} |\nabla \log(1+n(t))|^2 \Psi_R dx \\ + \int_{\mathbb{R}^2} |\nabla u|^2 \Psi_R dx = \int_{\mathbb{R}^2} n(t) \log(1+n(t)) \Psi_R dx + F_1(t) + F_2(t) + F_3(t), \end{aligned}$$

where

$$\begin{aligned} F_1(t) &= \int_{\mathbb{R}^2} [(1+n(t)) \log(1+n(t)) - n(t)] \Delta \Psi_R dx \\ &\quad - \int_{\mathbb{R}^2} n(t) c_m(t) \nabla c_m(t) \cdot \nabla \Psi_R dx + \int_{\mathbb{R}^2} c_m(t) \nabla n(t) \cdot \nabla \Psi_R dx \\ &\quad + \int_{\mathbb{R}^2} [n(t) \log(1+n(t)) - \log(1+n(t))] \nabla c_m(t) \cdot \nabla \Psi_R dx, \\ F_2(t) &= - \int_{\mathbb{R}^2} \partial_t c_m(t) \nabla c_m(t) \cdot \nabla \Psi_R dx - \frac{1}{4} \frac{d}{dt} \int_{\mathbb{R}^2} |c_m(t)|^2 \Delta \Psi_R dx, \end{aligned}$$

and

$$\begin{aligned} F_3(t) &= \int_{\mathbb{R}^2} [(1+n) \log(1+n) - n] u \cdot \nabla \Psi_R dx + \frac{1}{2} \int_{\mathbb{R}^2} |u|^2 u \cdot \nabla \Psi_R dx - \int_{\mathbb{R}^2} u \cdot \nabla P \Psi_R dx \\ &\quad - \int_{\mathbb{R}^2} (u \cdot \nabla) u \cdot \nabla \Psi_R dx - \int_{\mathbb{R}^2} n c_m u \cdot \nabla \Psi_R dx. \end{aligned} \tag{4.13}$$

Proof. The proof is identical to that of [68, Lemma 3.13] with an addition term $F_3(t)$ generated by the coupling of the velocity field u . \square

Lemma 4.14. For $F_1(t)$ in Lemma 4.13, it holds that

$$\sup_{0 < t < T} |F_1(t)| < \infty.$$

Proof. The proof is identical to that of [68, Lemma 3.14]. \square

Lemma 4.15. For $F_2(t)$ in Lemma 4.13, it holds that

$$\sup_{0 < t < T} \left| \int_0^t F_2(s) ds \right| < \infty.$$

Proof. We define $\tilde{\Psi}_R$ as in the proof of [68, Lemma 3.15]. Indeed, $\tilde{\Psi}_R(x) := \Psi_R(x)J_R(x)$, where $J_R \in C_0^\infty(\mathbb{R}^2)$, $J_R(x) \in [0, 1]$, $J_R(x) = 1$ for $R \leq |x| \leq 2R$,

$$\text{supp } J_R \subset B_{3R}(0) \setminus \overline{B_{R/2}(0)},$$

and

$$|\nabla J_R(x)| \leq CR^{-1}, \quad |\Delta J_R(x)| \leq CR^{-2}.$$

The support of $\tilde{\Psi}_R$ is contained in A_R , where A_R is an annulus defined in (4.7). In terms of $\tilde{\Psi}_R$, F_2 can be rewritten as

$$\begin{aligned} F_2(t) &= \frac{1}{4} \frac{d}{dt} \left\{ \int_{\mathbb{R}^2} c_m^2 \Delta \tilde{\Psi}_R dx - 2|\nabla c_m|^2 \tilde{\Psi}_R dx \right\} \\ &\quad + \int_{\mathbb{R}^2} [\Delta n - \nabla n \cdot (\nabla c_m + u) + n^2] c_m \tilde{\Psi}_R dx \\ &=: F_{21}(t) + F_{22}(t), \end{aligned}$$

where $F_{21}(t)$ is bounded as shown in the proof of [68, Lemma 3.15]. For $F_{22}(t)$, we rewrite it as

$$F_{22}(t) = -2 \int_{\mathbb{R}^2} \nabla n \cdot (\nabla c_m + u) \tilde{\Psi}_R dx - \int_{\mathbb{R}^2} \nabla n \cdot \nabla \tilde{\Psi}_R c_m dx + \int_{\mathbb{R}^2} n^2 c_m \tilde{\Psi}_R dx.$$

Since the integrands in $F_{22}(t)$ are bounded on $(0, T) \times \tilde{A}_R$ by (i) and (ii) of Proposition 4.12, we also have

$$\sup_{0 < t < T} |F_{22}(t)| < \infty.$$

This completes the proof of Lemma 4.15. \square

Lemma 4.16. *For $F_3(t)$ in Lemma 4.13, it holds that*

$$\sup_{0 < t < T} \left| \int_0^t F_3(s) ds \right| < \infty.$$

Proof. Note that

$$- \int_{\mathbb{R}^2} u \cdot \nabla P \Psi_R dx \leq \frac{1}{2} \|u(t)\|_2^2 + \frac{1}{2} \|\nabla P(t)\|_2^2 \in L^1(0, T)$$

by Remark 3.4, and all the other terms in (4.13) containing $\nabla \Psi_R$ are bounded by (i) and (ii) of Proposition 4.12. Therefore Lemma 4.16 is established. \square

The following proposition of interior estimate and the exterior estimate in Proposition 4.2 give the boundedness of the full modified entropy, which completes the proof of Proposition 1.4.

Proposition 4.17. *Assume $\int_{\mathbb{R}^2} n_0 dx \leq 8\pi$. Then*

$$\sup_{0 < t < T} \int_{|x| \leq 4\tilde{R}_0} (1 + n(t)) \log(1 + n(t)) dx < \infty,$$

where \tilde{R}_0 is the one determined in Proposition 4.12.

Proof. Denote $R = 4\tilde{R}_0$. It follows from Lemma 4.13 that

$$\frac{d}{dt} H_{\text{int}}(t; R) \leq \int_{\mathbb{R}^2} (1 + n(t)) \log(1 + n(t)) \Psi_R dx + F_1(t) + F_2(t) + F_3(t).$$

For $0 < a < 1$,

$$\frac{1}{2}nc_m \leq (1-a)n \frac{|c_m|}{2(1-a)} \leq (1-a)(1+n) \log(1+n) + (1-a) \exp \left\{ \frac{|c_m|}{2(1-a)} \right\},$$

from which it follows that

$$(1+n) \log(1+n) \leq \frac{1}{a} \left((1+n) \log(1+n) - \frac{1}{2}nc_m \right) + \frac{1-a}{a} \exp \left\{ \frac{|c_m|}{2(1-a)} \right\}.$$

Then

$$\begin{aligned} & \int_{\mathbb{R}^2} (1+n(t)) \log(1+n(t)) \Psi_R dx \\ & \leq \frac{1}{a} H_{\text{int}}(t; R) + \frac{1}{a} \int_{\mathbb{R}^2} n(t) \Psi_R dx + \frac{1-a}{a} \int_{\mathbb{R}^2} \exp \left\{ \frac{|c_m|}{2(1-a)} \right\} \Psi_R dx \\ & \leq \frac{1}{a} H_{\text{int}}(t; R) + \frac{1}{a} \|n_0\|_1 + F_4(t), \end{aligned} \quad (4.14)$$

where

$$F_4(t) = \frac{1-a}{a} \int_{\mathbb{R}^2} \exp \left\{ \frac{|c_m|}{2(1-a)} \right\} \Psi_R dx.$$

Therefore,

$$\frac{d}{dt} H_{\text{int}}(t; R) \leq \frac{1}{a} H_{\text{int}}(t; R) + F(t), \quad (4.15)$$

where

$$F(t) = F_1(t) + F_2(t) + F_3(t) + F_4(t) + \frac{1}{a} \|n_0\|_1.$$

We claim

$$\sup_{0 < t < T} F_4(t) < \infty. \quad (4.16)$$

To prove this claim, by (iii) of Proposition 4.12 there exist $x_0 \in \mathbb{R}^2$, $\varepsilon_0 > 0$, $\delta > 0$ with $R + \varepsilon_0 < |x_0| < 2R$ such that

$$n(x, t) \geq \delta \quad \text{for } 0 \leq t < T, \quad |x - x_0| \leq \varepsilon_0. \quad (4.17)$$

By (ii) of Proposition 4.12,

$$C := \sup_{0 < t < T} \|c_m(T)\|_{L^\infty(R \leq |x| \leq 2R)} < \infty.$$

The function $c_m(t)$ satisfies

$$-\Delta \left(\frac{c_m(t)}{2(1-a)} \right) = \frac{n(t)}{2(1-a)}, \quad |x| < R$$

and $\sup_{|x|=R} |c_m(t)| \leq C$ ($0 < t < T$). By (4.17), we have

$$\begin{aligned} \int_{|x| \leq R} n(t) dx & \leq \int_{\mathbb{R}^2} n(t) dx - \int_{|x-x_0| \leq \varepsilon_0} n(t) dx \\ & \leq 8\pi - \delta\pi\varepsilon_0^2 = \pi(8 - \delta\varepsilon_0^2) \end{aligned}$$

and hence

$$\int_{|x| \leq R} \frac{n(t)}{2(1-a)} dx \leq \frac{\pi}{2(1-a)} (8 - \delta\varepsilon_0^2).$$

We choose $0 < a < 1$ such that

$$\frac{\pi}{2(1-a)} (8 - \delta\varepsilon_0^2) < 4\pi$$

and apply Lemma 2.7 to get that for $0 < t < T$,

$$\begin{aligned} \int_{|x|<R} \exp \left\{ \frac{|c_m(t)|}{2(1-a)} \right\} dx &\leq \frac{32\pi(1-a)R^2}{\delta\varepsilon_0^2 - 8a} \exp \left\{ \sup_{|x|=R} \frac{|c_m(t)|}{2(1-a)} \right\} \\ &\leq \frac{32\pi(1-a)R^2}{\delta\varepsilon_0^2 - 8a} \exp \left\{ \frac{C}{2(1-a)} \right\}. \end{aligned}$$

Hence

$$\int_{\mathbb{R}^2} \exp \left\{ \frac{|c_m(t)|}{2(1-a)} \right\} \Psi_R dx \leq \left(\frac{32\pi(1-a)R^2}{\delta\varepsilon_0^2 - 8a} + 3\pi R^2 \right) \exp \left\{ \frac{C}{2(1-a)} \right\},$$

which implies (4.16).

By (4.15), we have

$$H_{\text{int}}(t; R) \leq H_{\text{int}}(0; R) + \frac{1}{a} \int_0^t H_{\text{int}}(s; R) ds + \sup_{0 < t < T} \left| \int_0^t F(s) ds \right|, \quad 0 < t < T$$

and by Lemma 4.14, Lemma 4.15, Lemma 4.16, and (4.16),

$$\sup_{0 < t < T} \left| \int_0^t F(s) ds \right| < \infty.$$

Applying the Gronwall inequality, we deduce that

$$\sup_{0 < t < T} H_{\text{int}}(t; R) < \infty.$$

Therefore, by this estimate, (4.14) and (4.16), we obtain

$$\sup_{0 < t < T} \int_{\mathbb{R}^2} (1 + n(t)) \log(1 + n(t)) \Psi_R dx < \infty,$$

which completes the proof of Proposition 4.17. \square

5. GLOBAL EXISTENCE OF THE PATLAK–KELLER–SEGEL–NAVIER–STOKES SYSTEM (1.1)

In this section, we prove our main result on global-in-time existence, Theorem 1.3.

Proposition 5.1. *Let (n_0, u_0) satisfy (1.3) and (1.4) with $M := \|n_0\|_1 = 8\pi$. For a local-in-time mild solution (n, u) on $[0, T)$ with initial data (n_0, u_0) given in Theorem 3.2. If $T < \infty$, it holds that for any $t_0 \in (0, T)$*

$$\sup_{t \in (t_0, T)} (\|n(t)\|_\infty + \|u(t)\|_\infty) < \infty.$$

Proof. The proof is divided into five steps.

Step 1. By testing (1.1)₁ with n and using the inequality (2.4), we get for $t \in (t_0, T)$, where $t_0 \in (0, T)$ is fixed, that

$$\begin{aligned} &\frac{1}{2} \|n(t)\|_2^2 + \int_{t_0}^t \|\nabla n(\tau)\|_2^2 d\tau \\ &= \frac{1}{2} \|n(t_0)\|_2^2 + \int_{t_0}^t \|n(\tau)\|_3^3 d\tau \\ &\leq \frac{1}{2} \|n(t_0)\|_2^2 + \varepsilon \left(\sup_{t \in (t_0, T)} \|(1 + n(t)) \log(1 + n(t))\|_1 \right) \int_{t_0}^t \|\nabla n(\tau)\|_2^2 d\tau + CMt \end{aligned}$$

since $\nabla \cdot u = 0$. It follows from Proposition 1.4 that $\sup_{t \in (t_0, T)} \|(1 + n(t)) \log(1 + n(t))\|_1 < \infty$. Thus we can choose $\varepsilon > 0$ sufficiently small so that for any $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|n(t)\|_2 < \infty \quad (5.1)$$

if $T < \infty$.

Step 2. Multiplying (1.1)₁ with n^{p-1} , we have

$$\frac{d}{dt} \left(\frac{n^p}{p} \right) + u \cdot \nabla \left(\frac{n^p}{p} \right) = \nabla \cdot (n^{p-1} \nabla n) - \frac{4(p-1)}{p^2} |\nabla(n^{p/2})|^2 - \nabla \cdot (n \nabla c) n^{p-1}.$$

Integrating over \mathbb{R}^2 , we obtain

$$\frac{1}{p} \frac{d}{dt} \|n(t)\|_p^p = -\frac{4(p-1)}{p^2} \int |\nabla(n^{p/2})|^2 + \frac{p-1}{p} \int n^{p+1}.$$

Taking $p = 4$,

$$\frac{1}{4} \frac{d}{dt} \|n(t)\|_4^4 = -\frac{3}{4} \int |\nabla(n^2)|^2 + \frac{3}{4} \int n^5.$$

By Gagliardo-Nirenberg-Sobolev inequality and Young's inequality, we have

$$\begin{aligned} \|n\|_5^5 &= \|n^2\|_{5/2}^{5/2} \leq C \|n^2\|_1 \|\nabla(n^2)\|_2^{3/2} = C \|n\|_2^2 \|\nabla(n^2)\|_2^{3/2} \\ &\leq \varepsilon \|\nabla(n^2)\|_2^2 + C \|n\|_2^8. \end{aligned}$$

Choosing $\varepsilon > 0$ sufficiently small, we have $\frac{d}{dt} \|n(t)\|_4^4 \leq C \|n\|_2^8$, and thus for $t \in (t_0, T)$, where $t_0 \in (0, T)$ is fixed, that

$$\|n(t)\|_4^4 \leq \|n(t_0)\|_4^4 + C \left(\sup_{s \in (t_0, T)} \|n(s)\|_2^8 \right) t.$$

Therefore, by (5.1), we get for any $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|n(t)\|_4 < \infty \quad (5.2)$$

if $T < \infty$.

Step 3. Since $\nabla \cdot u = 0$, $\|\nabla u\|_2 = \|\nabla \times u\|_2 = \|\omega\|_2$, where $\omega = \nabla \times u$. The vorticity equation of (1.1)₃ reads

$$\partial_t \omega + (u \cdot \nabla) \omega = \Delta \omega + \nabla \times (n \nabla c),$$

which has the following energy equality:

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|\omega(t)\|_2^2 + \|\nabla \omega(t)\|_2^2 &= \int_{\mathbb{R}^2} \nabla \times (n \nabla c) \omega \, dx \\ &= - \int_{\mathbb{R}^2} (n \nabla c) \cdot (\nabla^\perp \omega) \, dx, \end{aligned}$$

where we integrate by parts in the last equality. Thus, for $t \in (t_0, T)$, where $t_0 \in (0, T)$ is fixed,

$$\begin{aligned} \frac{1}{2} \|\omega(t)\|_2^2 + \int_{t_0}^t \|\nabla \omega(\tau)\|_2^2 \, d\tau &= \frac{1}{2} \|\omega(t_0)\|_2^2 - \int_{t_0}^t \int_{\mathbb{R}^2} (n \nabla c) \cdot (\nabla^\perp \omega)(\tau) \, dx \, d\tau \\ &\leq \frac{1}{2} \|\omega(t_0)\|_2^2 + \int_{t_0}^t \|(n \nabla c)(\tau)\|_2 \|\nabla^\perp \omega(\tau)\|_2 \, d\tau \\ &\leq \frac{1}{2} \|\omega(t_0)\|_2^2 + \int_{t_0}^t \left(\frac{\|(n \nabla c)(\tau)\|_2^2}{2} + \frac{\|\nabla^\perp \omega(\tau)\|_2^2}{2} \right) \, d\tau. \end{aligned}$$

By (5.1) and (5.2),

$$\begin{aligned} \sup_{t \in (t_0, T)} \|n \nabla c\|_2 &\leq \sup_{t \in (t_0, T)} \|n\|_4 \|\nabla c\|_4 \lesssim \sup_{t \in (t_0, T)} \|n\|_4 \|n\|_{4/3} \\ &\leq \sup_{t \in (t_0, T)} \|n\|_4 \|n\|_1^{1/2} \|n\|_2^{1/2} \leq C \end{aligned}$$

and $\|\nabla^\perp \omega\|_2 \leq \|\nabla \omega\|_2$. We have

$$\|\omega(t)\|_2^2 + \frac{1}{2} \int_{t_0}^t \|\nabla \omega(\tau)\|_2^2 d\tau \leq \|\omega(t_0)\|_2^2 + \frac{C^2}{2} T.$$

Therefore, we conclude that for any $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|\nabla u(t)\|_2 < \infty \quad (5.3)$$

if $T < \infty$.

Step 4. By (5.3) and the boundedness $\|u\|_2$ in Lemma 3.3 where the bound is independent of time since $M = 8\pi$, we have for $1 \leq q < \infty$ and $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|u(t)\|_q \lesssim \sup_{t \in (t_0, T)} \|u(t)\|_2^{8/q} \|\nabla u(t)\|_2^{8/q} \lesssim C \quad (5.4)$$

by the Sobolev's embedding theorem.

For $\tau \in (0, T - t_0)$, where $t_0 \in (0, T)$ is fixed,

$$u(t_0 + \tau) = e^{t_0 \Delta} u(\tau) - \int_0^{t_0} e^{(t_0-s)\Delta} \mathbf{P} \nabla \cdot [\nabla c(s + \tau) \otimes \nabla c(s + \tau) + u(s + \tau) \otimes u(s + \tau)] ds.$$

Then

$$\begin{aligned} \|u(t_0 + \tau)\|_\infty &\leq \|e^{t_0 \Delta} u(\tau)\|_\infty + C \int_0^{t_0} (t_0 - s)^{-\frac{1}{2} - \frac{1}{3}} \left(\|\nabla c(s + \tau)\|_6^2 + \|u(s + \tau)\|_6^2 \right) ds \\ &\lesssim t_0^{-\frac{1}{2}} \|u_0\|_2 + t_0^{\frac{1}{6}} \sup_{s \in (0, t_0)} \left(\|n(s + \tau)\|_{3/2}^2 + \|u(s + \tau)\|_6^2 \right) \\ &\lesssim t_0^{-\frac{1}{2}} \|u_0\|_2 + t_0^{\frac{1}{6}} \sup_{\sigma \in (\tau, T)} \left(\|n(\sigma)\|_1^{2/3} \|n(\sigma)\|_2^{4/3} + \|u(\sigma)\|_6^2 \right) \\ &\leq C \end{aligned}$$

by (5.1) and (5.4). Note that the bound is independent of τ since $\|u(t_0 + \tau)\|_\infty$ is bounded as $\tau \rightarrow 0_+$. Thus, we conclude for any $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|u(t)\|_\infty < \infty \quad (5.5)$$

if $T < \infty$.

Step 5. From Lemma 2.8, we have for any $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|\nabla c(t)\|_\infty \lesssim \sup_{t \in (t_0, T)} \|n(t)\|_1^{\frac{1}{3}} \|n(t)\|_4^{\frac{2}{3}} \leq C \quad (5.6)$$

by (5.2). For $\tau \in (0, T - t_0)$, where $t_0 \in (0, T)$ is fixed,

$$n(t_0 + \tau) = e^{t_0 \Delta} n(\tau) - \int_0^{t_0} e^{(t_0-s)\Delta} \nabla \cdot [n(s + \tau) \nabla c(s + \tau) + n(s + \tau) u(s + \tau)] ds.$$

Then

$$\begin{aligned}
& \|n(t_0 + \tau)\|_\infty \\
& \leq \|e^{t_0 \Delta} n(\tau)\|_\infty + C \int_0^{t_0} (t_0 - s)^{-\frac{1}{2} - \frac{1}{4}} (\|n(s + \tau) \nabla c(s + \tau)\|_4 + \|n(s + \tau) u(s + \tau)\|_4) ds \\
& \lesssim t_0^{-1} \|n(\tau)\|_1 + t_0^{\frac{1}{4}} \left[\sup_{\sigma \in (\tau, T)} (\|\nabla c(s + \tau)\|_\infty + \|u(s + \tau)\|_\infty) \right] \sup_{\sigma \in (\tau, T)} \|n(s + \tau)\|_4 \\
& \leq C
\end{aligned}$$

by (5.6), (5.5) and (5.2). Note that the bound is independent of τ since $\|n(t_0 + \tau)\|_\infty$ is bounded as $\tau \rightarrow 0_+$. Thus, we conclude for any $t_0 \in (0, T)$

$$\sup_{t \in (t_0, T)} \|n(t)\|_\infty < \infty \quad (5.7)$$

if $T < \infty$. The proposition follows from (5.5) and (5.7). \square

Proof of Theorem 1.3. Let T_m be the maximal existence time of (n, u) , and suppose $T_m < \infty$ by contradiction.

For the subcritical case $M < 8\pi$. Let $T < T_m$ be fixed. From (iv) of Theorem 3.2, $n(T)$ and $u(T)$ belong to $H^2(\mathbb{R}^2)$. According to [43, Theorem 1], solutions exist globally in time, which contradicts the definition of T_m . Moreover, (1.6) is a consequence of [43, (1.3)] and a use of Sobolev's embedding.

For the critical case $M = 8\pi$, we first claim that $n(t) \rightarrow \tilde{n}_0$ in $L^{4/3}$ and $u(t) \rightarrow \tilde{u}_0$ in L^2 as $t \rightarrow T_m -$ for some $(\tilde{n}_0, \tilde{u}_0) \in L^{4/3} \times L^2_\sigma$. Indeed, for fixed $t_0 \in (0, T_m)$, since

$$\begin{aligned}
\nabla^k n(t) &= D^k e^{(t-t_0/2)\Delta} n(t_0/2) - \int_{t_0}^t D^k \nabla e^{(t-s)\Delta} (n(s) \nabla c(s) + n(s) u(s)) ds \\
&=: \nabla^k e^{(t-t_0/2)\Delta} n(t_0/2) + I,
\end{aligned}$$

where, by [66, (3.1)],

$$\left\| D^k e^{(t-t_0/2)\Delta} n(t_0/2) \right\|_{4/3} \leq C t_0^{-k/2} \|n(t_0/2)\|_{4/3}$$

and

$$\begin{aligned}
\|I\|_{4/3} &\lesssim C \int_{t_0}^t (t-s)^{-(1-\frac{3}{4})-\frac{k+1}{2}} \|n(s) (\nabla c(s) + u(s))\|_1 ds \\
&\lesssim C \int_{t_0}^t (t-s)^{-\frac{3}{4}-\frac{k}{2}} \|n(s)\|_{4/3} \left(\|n(s)\|_{4/3} + \|u(s)\|_4 \right) ds
\end{aligned}$$

in which $\|n\|_{4/3}$ and $\|u\|_4$ are uniformly bounded in time by Proposition 5.1, we obtain for $0 < k < 1/2$ that

$$\sup_{t_0 \leq t \leq T_m} \|D^k n(t)\|_{4/3} < \infty,$$

and thus, using [66, (3.2)],

$$\left\| e^{(t-\tau)\Delta} n(\tau) - n(\tau) \right\|_{4/3} \leq C (t-\tau)^{k/2} \|D^k n(\tau)\|_{4/3} \rightarrow 0$$

as $\tau, t \rightarrow T_m -$. Therefore, $n_0(t) \rightarrow \tilde{n}_0$ in $L^{4/3}$ as $t \rightarrow T_m -$. Similarly, since

$$\left\| e^{(t-\tau)\Delta} u(\tau) - u(\tau) \right\|_4 \leq C (t-\tau)^{3/4} \|\nabla u(\tau)\|_2$$

in which $\sup_{t \in (t_0, T)} \|\nabla u(t)\|_2 < \infty$ by (5.3), we also have $u_0 \rightarrow \tilde{u}_0$ in L^4 as $t \rightarrow T_m^-$. It is obvious that $\tilde{n}_0 \in L^1 \cap L^{4/3}$ and $\tilde{u}_0 \in L^2 \cap L^4$. By Theorem 3.2 and Remark 3.1, there exists a unique mild solution (\tilde{n}, \tilde{u}) on $[0, T_0)$, $T_0 > 0$, with initial data $(\tilde{n}_0, \tilde{u}_0)$, satisfying $\tilde{n} \in BC([0, T_0]; L^1 \cap L^{4/3})$ and $\tilde{u} \in BC([0, T_0]; L^2 \cap L^4)$. Define (\hat{n}, \hat{u}) by

$$\hat{n}(t) = \begin{cases} n(t), & 0 \leq t < T_m, \\ \tilde{n}(t - T_m), & T_m \leq t < T_m + T_0, \end{cases}$$

$$\hat{u}(t) = \begin{cases} u(t), & 0 \leq t < T_m, \\ \tilde{u}(t - T_m), & T_m \leq t < T_m + T_0. \end{cases}$$

Then (n, u) is a mild solution on $[0, T_m + T_0)$, contradicting the choice of T_m . This proves the global-in-time existence. Finally, the L^∞ -boundedness (1.6) follows from Proposition 5.1. \square

ACKNOWLEDGEMENTS

The research of C. Lai is partially supported by FYF (#6456) of Graduate and Postdoctoral Studies, UBC. The research of J. Wei is partially supported by the NSERC grant of Canada.

REFERENCES

- [1] Hyeong-Ohk Bae and Lorenzo Brandolese. On the effect of external forces on incompressible fluid motions at large distances. *Ann. Univ. Ferrara Sez. VII Sci. Mat.*, 55(2):225–238, 2009.
- [2] Hajer Bahouri, Mohamed Majdoub, and Nader Masmoudi. On the lack of compactness in the 2D critical Sobolev embedding. *J. Funct. Anal.*, 260(1):208–252, 2011.
- [3] William Beckner. Sharp Sobolev inequalities on the sphere and the Moser-Trudinger inequality. *Ann. of Math. (2)*, 138(1):213–242, 1993.
- [4] Jacob Bedrossian and Siming He. Suppression of blow-up in Patlak-Keller-Segel via shear flows. *SIAM J. Math. Anal.*, 49(6):4722–4766, 2017.
- [5] Jacob Bedrossian and Siming He. Erratum: Suppression of blow-up in Patlak-Keller-Segel via shear flows [MR3730537]. *SIAM J. Math. Anal.*, 50(6):6365–6372, 2018.
- [6] Jacob Bedrossian and Nader Masmoudi. Existence, uniqueness and Lipschitz dependence for Patlak-Keller-Segel and Navier-Stokes in \mathbb{R}^2 with measure-valued initial data. *Arch. Ration. Mech. Anal.*, 214(3):717–801, 2014.
- [7] Matania Ben-Artzi. Global solutions of two-dimensional Navier-Stokes and Euler equations. *Arch. Rational Mech. Anal.*, 128(4):329–358, 1994.
- [8] Piotr Biler. The Cauchy problem and self-similar solutions for a nonlinear parabolic equation. *Studia Math.*, 114(2):181–205, 1995.
- [9] Piotr Biler. Existence and nonexistence of solutions for a model of gravitational interaction of particles. III. *Colloq. Math.*, 68(2):229–239, 1995.
- [10] Piotr Biler. Local and global solvability of some parabolic systems modelling chemotaxis. *Adv. Math. Sci. Appl.*, 8(2):715–743, 1998.
- [11] Piotr Biler, Danielle Hilhorst, and Tadeusz Nadzieja. Existence and nonexistence of solutions for a model of gravitational interaction of particles. II. *Colloq. Math.*, 67(2):297–308, 1994.
- [12] Piotr Biler, Grzegorz Karch, Philippe Laurençot, and Tadeusz Nadzieja. The 8π -problem for radially symmetric solutions of a chemotaxis model in the plane. *Math. Methods Appl. Sci.*, 29(13):1563–1583, 2006.
- [13] Piotr Biler and Tadeusz Nadzieja. A nonlocal singular parabolic problem modelling gravitational interaction of particles. *Adv. Differential Equations*, 3(2):177–197, 1998.
- [14] Piotr Biler and Jacek Zienkiewicz. Existence of solutions for the Keller-Segel model of chemotaxis with measures as initial data. *Bull. Pol. Acad. Sci. Math.*, 63(1):41–51, 2015.
- [15] Adrien Blanchet. On the parabolic-elliptic Patlak-Keller-Segel system in dimension 2 and higher. In *Séminaire Laurent Schwartz—Équations aux dérivées partielles et applications. Année 2011–2012*, Sémin. Équ. Dériv. Partielles, pages Exp. No. VIII, 26. École Polytech., Palaiseau, 2013.
- [16] Adrien Blanchet, Eric A. Carlen, and José A. Carrillo. Functional inequalities, thick tails and asymptotics for the critical mass Patlak-Keller-Segel model. *J. Funct. Anal.*, 262(5):2142–2230, 2012.

- [17] Adrien Blanchet, José A. Carrillo, and Nader Masmoudi. Infinite time aggregation for the critical Patlak-Keller-Segel model in \mathbb{R}^2 . *Comm. Pure Appl. Math.*, 61(10):1449–1481, 2008.
- [18] Adrien Blanchet, Jean Dolbeault, Miguel Escobedo, and Javier Fernández. Asymptotic behaviour for small mass in the two-dimensional parabolic-elliptic Keller-Segel model. *J. Math. Anal. Appl.*, 361(2):533–542, 2010.
- [19] Adrien Blanchet, Jean Dolbeault, and Benoît Perthame. Two-dimensional Keller-Segel model: optimal critical mass and qualitative properties of the solutions. *Electron. J. Differential Equations*, pages No. 44, 32, 2006.
- [20] Lorenzo Brandolese and Maria E. Schonbek. Large time behavior of the Navier-Stokes flow. In *Handbook of mathematical analysis in mechanics of viscous fluids*, pages 579–645. Springer, Cham, 2018.
- [21] Haïm Brezis. Remarks on the preceding paper by M. Ben-Artzi: “Global solutions of two-dimensional Navier-Stokes and Euler equations” [Arch. Rational Mech. Anal. **128** (1994), no. 4, 329–358; MR1308857 (96h:35148)]. *Arch. Rational Mech. Anal.*, 128(4):359–360, 1994.
- [22] Juan F. Campos and Jean Dolbeault. Asymptotic estimates for the parabolic-elliptic Keller-Segel model in the plane. *Comm. Partial Differential Equations*, 39(5):806–841, 2014.
- [23] E. Carlen and M. Loss. Competing symmetries, the logarithmic HLS inequality and Onofri’s inequality on S^n . *Geom. Funct. Anal.*, 2(1):90–104, 1992.
- [24] Jose A. Carrillo, Li Chen, Jian-Guo Liu, and Jinhuan Wang. A note on the subcritical two dimensional Keller-Segel system. *Acta Appl. Math.*, 119:43–55, 2012.
- [25] Pierre-Henri Chavanis and Clément Sire. Virial theorem and dynamical evolution of self-gravitating Brownian particles in an unbounded domain. I. Overdamped models. *Phys. Rev. E (3)*, 73(6):066103, 16, 2006.
- [26] Charles Collot, Tej-Eddine Ghouil, Nader Masmoudi, and Van Tien Nguyen. Refined description and stability for singular solutions of the 2d keller-segel system. *arXiv preprint arXiv:1912.00721*, 2019.
- [27] Charles Collot, Tej-Eddine Ghouil, Nader Masmoudi, and Van Tien Nguyen. Refined description and stability for singular solutions of the 2d keller-segel system. *arXiv preprint arXiv:1912.00721*, 2019.
- [28] Peter Constantin and Mihaela Ignatova. On the Nernst-Planck-Navier-Stokes system. *Arch. Ration. Mech. Anal.*, 232(3):1379–1428, 2019.
- [29] Juan Davila, Manuel del Pino, Jean Dolbeault, Monica Musso, and Juncheng Wei. Infinite time blow-up in the patlak-keller-segel system: existence and stability. *arXiv preprint arXiv:1911.12417*, 2019.
- [30] Jean-Marc Delort. Existence de nappes de tourbillon en dimension deux. *J. Amer. Math. Soc.*, 4(3):553–586, 1991.
- [31] Jesus Ildefonso Diaz, Toshitaka Nagai, and Jean-Michel Rakotoson. Symmetrization techniques on unbounded domains: application to a chemotaxis system on \mathbf{R}^N . *J. Differential Equations*, 145(1):156–183, 1998.
- [32] Jean Dolbeault and Benoît Perthame. Optimal critical mass in the two-dimensional Keller-Segel model in \mathbb{R}^2 . *C. R. Math. Acad. Sci. Paris*, 339(9):611–616, 2004.
- [33] Jean Dolbeault and Christian Schmeiser. The two-dimensional Keller-Segel model after blow-up. *Discrete Contin. Dyn. Syst.*, 25(1):109–121, 2009.
- [34] Herbert Gajewski and Klaus Zacharias. Global behaviour of a reaction-diffusion system modelling chemotaxis. *Math. Nachr.*, 195:77–114, 1998.
- [35] G. P. Galdi. *An introduction to the mathematical theory of the Navier-Stokes equations*. Springer Monographs in Mathematics. Springer, New York, second edition, 2011. Steady-state problems.
- [36] Isabelle Gallagher and Thierry Gallay. Uniqueness for the two-dimensional Navier-Stokes equation with a measure as initial vorticity. *Math. Ann.*, 332(2):287–327, 2005.
- [37] Thierry Gallay and C. Eugene Wayne. Invariant manifolds and the long-time asymptotics of the Navier-Stokes and vorticity equations on \mathbf{R}^2 . *Arch. Ration. Mech. Anal.*, 163(3):209–258, 2002.
- [38] Thierry Gallay and C. Eugene Wayne. Global stability of vortex solutions of the two-dimensional Navier-Stokes equation. *Comm. Math. Phys.*, 255(1):97–129, 2005.
- [39] Tej-Eddine Ghouil and Nader Masmoudi. Minimal mass blowup solutions for the Patlak-Keller-Segel equation. *Comm. Pure Appl. Math.*, 71(10):1957–2015, 2018.
- [40] Y. Giga, S. Matsui, and O. Sawada. Global existence of two-dimensional Navier-Stokes flow with nondecaying initial velocity. *J. Math. Fluid Mech.*, 3(3):302–315, 2001.
- [41] Yoshikazu Giga, Tetsuro Miyakawa, and Hirofumi Osada. Two-dimensional Navier-Stokes flow with measures as initial vorticity. *Arch. Rational Mech. Anal.*, 104(3):223–250, 1988.
- [42] Yoshikazu Giga and Hermann Sohr. Abstract L^p estimates for the Cauchy problem with applications to the Navier-Stokes equations in exterior domains. *J. Funct. Anal.*, 102(1):72–94, 1991.
- [43] Yishu Gong and Siming He. On the 8π -subcritical mass threshold of a Patlak-Keller-Segel-Navier-Stokes system. *arXiv preprint arXiv:2003.12174*, 2020.

- [44] Siming He. Suppression of blow-up in parabolic-parabolic Patlak-Keller-Segel via strictly monotone shear flows. *Nonlinearity*, 31(8):3651–3688, 2018.
- [45] Miguel A. Herrero and Juan J. L. Velázquez. Chemotactic collapse for the Keller-Segel model. *J. Math. Biol.*, 35(2):177–194, 1996.
- [46] Miguel A. Herrero and Juan J. L. Velázquez. Singularity patterns in a chemotaxis model. *Math. Ann.*, 306(3):583–623, 1996.
- [47] Miguel A. Herrero and Juan J. L. Velázquez. A blow-up mechanism for a chemotaxis model. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 24(4):633–683 (1998), 1997.
- [48] Eberhard Hopf. Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen. *Math. Nachr.*, 4:213–231, 1951.
- [49] Dirk Horstmann. From 1970 until present: the Keller-Segel model in chemotaxis and its consequences. I. *Jahresber. Deutsch. Math.-Verein.*, 105(3):103–165, 2003.
- [50] Dirk Horstmann. From 1970 until present: the Keller-Segel model in chemotaxis and its consequences. II. *Jahresber. Deutsch. Math.-Verein.*, 106(2):51–69, 2004.
- [51] W. Jäger and S. Luckhaus. On explosions of solutions to a system of partial differential equations modelling chemotaxis. *Trans. Amer. Math. Soc.*, 329(2):819–824, 1992.
- [52] Tosio Kato. Strong L^p -solutions of the Navier-Stokes equation in \mathbf{R}^m , with applications to weak solutions. *Math. Z.*, 187(4):471–480, 1984.
- [53] Tosio Kato. The Navier-Stokes equation for an incompressible fluid in \mathbf{R}^2 with a measure as the initial vorticity. *Differential Integral Equations*, 7(3-4):949–966, 1994.
- [54] Evelyn F. Keller and Lee A. Segel. Initiation of slime mold aggregation viewed as an instability. *J. Theoret. Biol.*, 26(3):399–415, 1970.
- [55] Alexander Kiselev and Xiaoqian Xu. Suppression of chemotactic explosion by mixing. *Arch. Ration. Mech. Anal.*, 222(2):1077–1112, 2016.
- [56] Chen-Chih Lai, Juncheng Wei, and Yifu Zhou. Infinite time blow-up for the 2D PKS-NS system with critical mass. *preprint*, 2021.
- [57] P. G. Lemarié-Rieusset. *Recent developments in the Navier-Stokes problem*, volume 431 of *Chapman & Hall/CRC Research Notes in Mathematics*. Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [58] Jean Leray. Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.*, 63(1):193–248, 1934.
- [59] Pierre-Louis Lions. *Mathematical topics in fluid mechanics. Vol. 1*, volume 3 of *Oxford Lecture Series in Mathematics and its Applications*. The Clarendon Press, Oxford University Press, New York, 1996. Incompressible models, Oxford Science Publications.
- [60] Julián López-Gómez, Toshitaka Nagai, and Tetsuya Yamada. The basin of attraction of the steady-states for a chemotaxis model in \mathbb{R}^2 with critical mass. *Arch. Ration. Mech. Anal.*, 207(1):159–184, 2013.
- [61] S. Luckhaus, Y. Sugiyama, and J. J. L. Velázquez. Measure valued solutions of the 2D Keller-Segel system. *Arch. Ration. Mech. Anal.*, 206(1):31–80, 2012.
- [62] Noriko Mizoguchi. Global existence for the Cauchy problem of the parabolic-parabolic Keller-Segel system on the plane. *Calc. Var. Partial Differential Equations*, 48(3-4):491–505, 2013.
- [63] Noriko Mizoguchi. Refined asymptotic behavior of blowup solutions to a simplified chemotaxis system. *Communications on Pure and Applied Mathematics*, 2020.
- [64] Toshitaka Nagai. Blow-up of radially symmetric solutions to a chemotaxis system. *Adv. Math. Sci. Appl.*, 5(2):581–601, 1995.
- [65] Toshitaka Nagai. Blowup of nonradial solutions to parabolic-elliptic systems modeling chemotaxis in two-dimensional domains. *J. Inequal. Appl.*, 6(1):37–55, 2001.
- [66] Toshitaka Nagai. Global existence and decay estimates of solutions to a parabolic-elliptic system of drift-diffusion type in \mathbb{R}^2 . *Differential Integral Equations*, 24(1-2):29–68, 2011.
- [67] Toshitaka Nagai and Takayoshi Ogawa. Brezis-Merle inequalities and application to the global existence of the Cauchy problem of the Keller-Segel system. *Commun. Contemp. Math.*, 13(5):795–812, 2011.
- [68] Toshitaka Nagai and Takayoshi Ogawa. Global existence of solutions to a parabolic-elliptic system of drift-diffusion type in \mathbf{R}^2 . *Funkcial. Ekvac.*, 59(1):67–112, 2016.
- [69] Toshitaka Nagai, Takasi Senba, and Kiyoshi Yoshida. Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis. *Funkcial. Ekvac.*, 40(3):411–433, 1997.
- [70] Vidyanand Nanjundiah. Chemotaxis, signal relaying and aggregation morphology. *Journal of Theoretical Biology*, 42(1):63–105, 1973.
- [71] Takayoshi Ogawa, Shubha V. Rajopadhye, and Maria E. Schonbek. Energy decay for a weak solution of the Navier-Stokes equation with slowly varying external forces. *J. Funct. Anal.*, 144(2):325–358, 1997.
- [72] Clifford S. Patlak. Random walk with persistence and external bias. *Bull. Math. Biophys.*, 15:311–338, 1953.

- [73] Frédéric Poupaud. Diagonal defect measures, adhesion dynamics and Euler equation. *Methods Appl. Anal.*, 9(4):533–561, 2002.
- [74] Pierre Raphaël and Rémi Schweyer. On the stability of critical chemotactic aggregation. *Math. Ann.*, 359(1-2):267–377, 2014.
- [75] Gregory Seregin. *Lecture notes on regularity theory for the Navier-Stokes equations*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2015.
- [76] Hermann Sohr. *The Navier-Stokes equations*. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2001. An elementary functional analytic approach.
- [77] Elias M. Stein. *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, volume 43 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1993. With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III.
- [78] Roger Temam. *Navier-Stokes equations*. AMS Chelsea Publishing, Providence, RI, 2001. Theory and numerical analysis, Reprint of the 1984 edition.
- [79] Neil S. Trudinger. On imbeddings into Orlicz spaces and some applications. *J. Math. Mech.*, 17:473–483, 1967.
- [80] Tai-Peng Tsai. *Lectures on Navier-Stokes equations*, volume 192 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2018.
- [81] J. J. L. Velázquez. Stability of some mechanisms of chemotactic aggregation. *SIAM J. Appl. Math.*, 62(5):1581–1633, 2002.
- [82] J. J. L. Velázquez. Point dynamics in a singular limit of the Keller-Segel model. I. Motion of the concentration regions. *SIAM J. Appl. Math.*, 64(4):1198–1223, 2004.
- [83] J. J. L. Velázquez. Point dynamics in a singular limit of the Keller-Segel model. I. Motion of the concentration regions. *SIAM J. Appl. Math.*, 64(4):1198–1223, 2004.
- [84] J. J. L. Velázquez. Point dynamics in a singular limit of the Keller-Segel model. II. Formation of the concentration regions. *SIAM J. Appl. Math.*, 64(4):1224–1248, 2004.
- [85] Dongyi Wei. Global well-posedness and blow-up for the 2-D Patlak-Keller-Segel equation. *J. Funct. Anal.*, 274(2):388–401, 2018.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, B.C., V6T 1Z2, CANADA
Email address: chenchih@math.ubc.ca

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, B.C., V6T 1Z2, CANADA
Email address: jcwei@math.ubc.ca

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, 3400 N. CHARLES STREET, BALTIMORE, MD 21218, USA
Email address: yzhou173@math.jhu.edu